

## Errata and Addenda to *Mathematical Constants*

STEVEN FINCH

April 12, 2007

At this point, there are more additions than errors to report...

**1.1. Pythagoras' Constant.** A geometric irrationality proof of  $\sqrt{2}$  appears in [1] and a curious recursion is studied in [2]. More references on radical denestings include [3, 4, 5, 6].

**1.2. The Golden Mean.** The cubic irrational  $\chi = 1.8392867552\dots$  is mentioned elsewhere in the literature with regard to iterative functions [7, 8, 9] (the four-numbers game is a special case of what are known as Ducci sequences), geometric constructions [10, 11] and numerical analysis [12]. Another reference on infinite radical expressions is [13]. See [14] for an interesting optimality property of the logarithmic spiral. A mean-value analog  $C$  of Viswanath's constant 1.13198824... (the latter applies for almost every random Fibonacci sequence) has been estimated by Makover & McGowan [15]; they deduce that  $1.12095 \leq C \leq 1.23375$ . The Fibonacci factorial constant  $c$  arises in [16] with regard to the asymptotics

$$\begin{aligned} -\frac{d}{ds} \sum_{n=1}^{\infty} \frac{1}{f_n^s} &\sim \frac{1}{\ln(\varphi)s^2} + \frac{1}{24} \left( 6 \ln(5) - 2 \ln(\varphi) - \frac{3 \ln(5)^2}{\ln(\varphi)} \right) + \ln(c) \\ &\sim \frac{1}{\ln(\varphi)s^2} + \ln(0.8992126807\dots) \end{aligned}$$

as  $s \rightarrow 0$ , which gives meaning to the “regularized product” of all Fibonacci numbers.

**1.3. The Natural Logarithmic Base.** A proof of the formula

$$\frac{e}{2} = \left(\frac{2}{1}\right)^{\frac{1}{2}} \cdot \left(\frac{2 \cdot 4}{3 \cdot 3}\right)^{\frac{1}{4}} \cdot \left(\frac{4 \cdot 6 \cdot 6 \cdot 8}{5 \cdot 5 \cdot 7 \cdot 7}\right)^{\frac{1}{8}} \cdots$$

appears in [17]; Hurwitzian continued fractions for  $e^{1/q}$  and  $e^{2/q}$  appear in [18, 19, 20, 21]. Define the following set of integer  $k$ -tuples

$$N_k = \left\{ (n_1, n_2, \dots, n_k) : \sum_{j=1}^k \frac{1}{n_j} = 1 \text{ and } 1 \leq n_1 < n_2 < \dots < n_k \right\}.$$

Martin [22] proved that

$$\min_{(n_1, n_2, \dots, n_k) \in N_k} n_k \sim \frac{e}{e-1} k$$

---

<sup>0</sup>Copyright © 2007 by Steven R. Finch. All rights reserved.

as  $k \rightarrow \infty$ , but it remains open whether

$$\max_{(n_1, n_2, \dots, n_k) \in N_k} n_1 \sim \frac{1}{e-1} k.$$

Croot [23] made some progress on the latter: He proved that  $n_1 \geq (1 + o(1))k/(e-1)$  for infinitely many values of  $k$ , and this bound is best possible. Also, define  $f_0(x) = x$  and, for each  $n > 0$ ,

$$f_n(x) = (1 + f_{n-1}(x) - f_{n-1}(0))^{\frac{1}{x}}.$$

This imitates the definition of  $e$ , in the sense that the exponent  $\rightarrow \infty$  and the base  $\rightarrow 1$  as  $x \rightarrow 0$ . We have  $f_1(0) = e = 2.718\dots$ ,

$$f_2(0) = \exp\left(-\frac{e}{2}\right) = 0.257\dots, \quad f_3(0) = \exp\left(\frac{11-3e}{24} \exp\left(1 - \frac{e}{2}\right)\right) = 1.086\dots$$

and  $f_4(0) = 0.921\dots$  (too complicated an expression to include here). Does a pattern develop here?

#### 1.4. Archimedes' Constant. Viète's product

$$\frac{2}{\pi} = \sqrt{\frac{1}{2}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}}} \cdots$$

has the following close cousin:

$$\frac{2}{L} = \sqrt{\frac{1}{2}} \cdot \sqrt{\frac{1}{2} + \frac{\frac{1}{2}}{\sqrt{\frac{1}{2}}}} \cdot \sqrt{\frac{1}{2} + \frac{\frac{1}{2}}{\sqrt{\frac{1}{2} + \frac{\frac{1}{2}}{\sqrt{\frac{1}{2}}}}}} \cdots$$

where  $L$  is the lemniscate constant (pages 420–423). Levin [24, 25] developed analogs of sine and cosine for the curve  $x^4 + y^4 = 1$  to prove the latter formula; he also noted that the area enclosed by  $x^4 + y^4 = 1$  is  $\sqrt{2}L$  and that

$$\frac{2\sqrt{3}}{\pi} = \left(\frac{1}{2} + \sqrt{\frac{1}{2}}\right) \cdot \left(\frac{1}{2} + \sqrt{\frac{1}{2} - \frac{1}{2}\sqrt{\frac{1}{2}}}\right) \cdot \left(\frac{1}{2} + \sqrt{\frac{1}{2} - \frac{1}{2}\sqrt{\frac{1}{2} - \frac{1}{2}\sqrt{\frac{1}{2}}}}\right) \cdots$$

Can the half-circumference of  $x^4 + y^4 = 1$  be written in terms of  $L$  as well? This question makes sense both in the usual 2-norm and in the 4-norm; call the half-circumference  $\pi_4$  for the latter. More generally, define  $\pi_p$  to be the half-circumference of the unit  $p$ -circle  $|x|^p + |y|^p = 1$ , where lengths are measured via the  $p$ -norm and  $1 \leq p < \infty$ . It turns out [26] that  $\pi = \pi_2$  is the minimum value of  $\pi_p$ . Additional

infinite radical expressions for  $\pi$  appear in [27, 28]; more on the Matiyasevich-Guy formula is covered in [29, 30, 31, 32, 33]; see [34] for a revised spigot algorithm for computing decimal digits of  $\pi$  and [35, 36] for more on BBP-type formulas.

**1.5. Euler-Mascheroni Constant.** De la Vallée Poussin's theorem was, in fact, anticipated by Dirichlet [37, 38]; it is a corollary of the formula for the limiting mean value of  $d(n)$  [39]. Vacca's series was anticipated by Nielsen [40] and Jacobsthal [41, 42]. An extension was found by Koecher [43]:

$$\gamma = \delta - \frac{1}{2} \sum_{k=2}^{\infty} \frac{(-1)^k}{(k-1)k(k+1)} \left\lfloor \frac{\ln(k)}{\ln(2)} \right\rfloor$$

where  $\delta = (1 + \alpha)/4 = 0.6516737881\dots$  and  $\alpha = \sum_{n=1}^{\infty} 1/(2^n - 1) = 1.6066951524\dots$  is one of the digital search tree constants. Glaisher [44] discovered a similar formula:

$$\gamma = \sum_{n=1}^{\infty} \frac{1}{3^n - 1} - 2 \sum_{k=1}^{\infty} \frac{1}{(3k-1)(3k)(3k+1)} \left\lfloor \frac{\ln(3k)}{\ln(3)} \right\rfloor$$

nearly eighty years earlier. The following series [45, 46, 47] suggest that  $\ln(4/\pi)$  is an “alternating Euler constant”:

$$\gamma = \sum_{k=1}^{\infty} \left( \frac{1}{k} - \ln \left( 1 + \frac{1}{k} \right) \right) = - \int_0^1 \int_0^1 \frac{1-x}{(1-xy) \ln(xy)} dx dy,$$

$$\ln \left( \frac{4}{\pi} \right) = \sum_{k=1}^{\infty} (-1)^{k-1} \left( \frac{1}{k} - \ln \left( 1 + \frac{1}{k} \right) \right) = - \int_0^1 \int_0^1 \frac{1-x}{(1+xy) \ln(xy)} dx dy$$

(see section 1.7 later for more). Sample criteria for the irrationality of  $\gamma$  appear in Sondow [48, 49, 50, 51, 52]. Long ago, Mahler attempted to prove that  $\gamma$  is transcendental; the closest he came to this was to prove the transcendentality of the constant [53, 54]

$$\frac{\pi Y_0(2)}{2J_0(2)} - \gamma$$

where  $J_0(x)$  and  $Y_0(x)$  are the zeroth Bessel functions of the first and second kinds. (Unfortunately the conclusion cannot be applied to the terms separately!) Diamond [55, 56] proved that, if

$$F_k(n) = \sum \frac{1}{\ln(\nu_1) \ln(\nu_2) \cdots \ln(\nu_k)}$$

where the (finite) sum is over all integer multiplicative compositions  $n = \nu_1 \nu_2 \cdots \nu_k$  and each  $\nu_j \geq 2$ , then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \left( 1 + \sum_{n=2}^N \sum_{k=1}^{\infty} \frac{F_k(n)}{k!} \right) = \exp(\gamma' - \gamma - \ln(\ln(2))) = 1.2429194164\dots$$

where  $\gamma' = 0.4281657248\dots$  is the analog of Euler's constant when  $1/x$  is replaced by  $1/(x \ln(x))$  (see Table 1.1). See [57] for a different generalization of  $\gamma$ .

**1.6. Apéry's Constant.** The famous alternating central binomial series for  $\zeta(3)$  dates back at least as far as 1890, appearing as a special case of a formula due to Markov [58, 59, 60]:

$$\sum_{n=0}^{\infty} \frac{1}{(x+n)^3} = \frac{1}{4} \sum_{n=0}^{\infty} \frac{(-1)^n (n!)^6}{(2n+1)!} \frac{2(x-1)^2 + 6(n+1)(x-1) + 5(n+1)^2}{[x(x+1)\cdots(x+n)]^4}.$$

Ramanujan [61, 62] discovered the series for  $\zeta(3)$  attributed to Grosswald. Plouffe [63] uncovered remarkable formulas for  $\pi^{2k+1}$  and  $\zeta(2k+1)$ , including

$$\begin{aligned} \pi &= 72 \sum_{n=1}^{\infty} \frac{1}{n(e^{\pi n} - 1)} - 96 \sum_{n=1}^{\infty} \frac{1}{n(e^{2\pi n} - 1)} + 24 \sum_{n=1}^{\infty} \frac{1}{n(e^{4\pi n} - 1)}, \\ \pi^3 &= 720 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{\pi n} - 1)} - 900 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{2\pi n} - 1)} + 180 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{4\pi n} - 1)}, \\ \pi^5 &= 7056 \sum_{n=1}^{\infty} \frac{1}{n^5(e^{\pi n} - 1)} - 6993 \sum_{n=1}^{\infty} \frac{1}{n^5(e^{2\pi n} - 1)} + 63 \sum_{n=1}^{\infty} \frac{1}{n^5(e^{4\pi n} - 1)}, \\ \zeta(3) &= 28 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{\pi n} - 1)} - 37 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{2\pi n} - 1)} + 7 \sum_{n=1}^{\infty} \frac{1}{n^3(e^{4\pi n} - 1)}, \\ \zeta(5) &= 24 \sum_{n=1}^{\infty} \frac{1}{n^5(e^{\pi n} - 1)} - \frac{259}{10} \sum_{n=1}^{\infty} \frac{1}{n^5(e^{2\pi n} - 1)} - \frac{1}{10} \sum_{n=1}^{\infty} \frac{1}{n^5(e^{4\pi n} - 1)}, \\ \zeta(7) &= \frac{304}{13} \sum_{n=1}^{\infty} \frac{1}{n^7(e^{\pi n} - 1)} - \frac{103}{4} \sum_{n=1}^{\infty} \frac{1}{n^7(e^{2\pi n} - 1)} + \frac{19}{52} \sum_{n=1}^{\infty} \frac{1}{n^7(e^{4\pi n} - 1)}. \end{aligned}$$

**1.7. Catalan's Constant.** Rivoal & Zudilin [64] proved that there exist infinitely many integers  $k$  for which  $\beta(2k)$  is irrational, and that at least one of the numbers  $\beta(2), \beta(4), \beta(6), \beta(8), \beta(10), \beta(12), \beta(14)$  is irrational. More double integrals (see section 1.5 earlier) include [65, 66, 67, 68]

$$\zeta(3) = -\frac{1}{2} \int_0^1 \int_0^1 \frac{\ln(xy) \, dx \, dy}{1-xy}, \quad G = \frac{1}{8} \int_0^1 \int_0^1 \frac{dx \, dy}{(1-xy)\sqrt{x(1-y)}}.$$

Zudilin [67] also found the continued fraction expansion

$$\frac{13}{2G} = 7 + \frac{1040}{|10699} + \frac{42322176}{|434871} + \frac{15215850000}{|4090123} + \dots$$

where the partial numerators and partial denominators are generated according to the polynomials  $(2n - 1)^4(2n)^4(20n^2 - 48n + 29)(20n^2 + 32n + 13)$  and  $3520n^6 + 5632n^5 + 2064n^4 - 384n^3 - 156n^2 + 16n + 7$ .

**1.8. Khintchine-Lévy Constants.** If  $x$  is a quadratic irrational, then its continued fraction expansion is periodic; hence  $\lim_{n \rightarrow \infty} M(n, x)$  is easily found and is algebraic. For example,  $\lim_{n \rightarrow \infty} M(n, \varphi) = 1$ , where  $\varphi$  is the Golden mean. We study the set  $\Sigma$  of values  $\lim_{n \rightarrow \infty} \ln(Q_n)/n$  taken over all quadratic irrationals  $x$  in [69]. Additional references include [70, 71, 72].

**1.9. Feigenbaum-Coulet-Tresser Constants.** Consider the unique solution of  $\varphi(x) = T_2[\varphi](x)$  as pictured in Figure 1.6. The Hausdorff dimension  $D$  of the Cantor set  $\{x_k\}_{k=1}^{\infty} \subseteq [-1, 1]$ , defined by  $x_1 = 1$  and  $x_{k+1} = \varphi(x_k)$ , is known to satisfy  $0.53763 < D < 0.53854$ . This set may be regarded as the simplest of all strange attractors [73, 74].

**1.11. Chaitin's Constant.** Ord & Kieu [75] gave a different Diophantine representation for  $\Omega$ ; apparently Chaitin's equation can be reduced to 2–3 pages in length [76]. A rough sense of the type of equations involved can be gained from [77]. Calude & Stay [78] suggested that the uncomputability of bits of  $\Omega$  can be recast as a uncertainty principle.

**2.1. Hardy-Littlewood Constants.** Green & Tao [79] recently proved that there are arbitrarily long arithmetic progressions of primes. In particular, the number of prime triples  $p_1 < p_2 < p_3 \leq x$  in arithmetic progression is

$$\sim \frac{C_{\text{twin}}}{2} \frac{x^2}{\ln(x)^3} = (0.3300809079\dots) \frac{x^2}{\ln(x)^3}$$

as  $x \rightarrow \infty$ , and the number of prime quadruples  $p_1 < p_2 < p_3 < p_4 \leq x$  in arithmetic progression is likewise

$$\sim \frac{D}{6} \frac{x^2}{\ln(x)^4} = (0.4763747659\dots) \frac{x^2}{\ln(x)^4}.$$

Fix  $\varepsilon > 0$ . Let  $N(x, k)$  denote the number of positive integers  $n \leq x$  with  $\Omega(n) = k$ , where  $k$  is allowed to grow with  $x$ . Nicolas [80] proved that

$$\lim_{x \rightarrow \infty} \frac{N(x, k)}{(x/2^k) \ln(x/2^k)} = \frac{1}{4C_{\text{twin}}} = \frac{1}{4} \prod_{p>2} \left( 1 + \frac{1}{p(p-2)} \right) = 0.3786950320\dots$$

under the assumption that  $(2 + \varepsilon) \ln(\ln(x)) \leq k \leq \ln(x)/\ln(2)$ . More relevant results appear in [81]; see also the next entry.

Let  $L(x)$  denote the number of positive odd integers  $n \leq x$  that can be expressed in the form  $2^l + p$ , where  $l$  is a positive integer and  $p$  is a prime. Then  $0.0868x <$

$L(x) \leq 0.49999991x$  for all sufficiently large  $x$ . The lower bound can be improved to  $0.2893x$  if the Hardy-Littlewood conjectures in sieve theory are true [82, 83, 84].

Let  $Q(x)$  denote the number of integers  $\leq x$  with prime factorizations  $p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$  satisfying  $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_r$ . Extending results of Hardy & Ramanujan [85], Richmond [86] deduced that

$$\ln(Q(x)) \sim \frac{2\pi}{\sqrt{3}} \left( \frac{\ln(x)}{\ln(\ln(x))} \right)^{1/2} \left( 1 - \frac{2\ln(\pi)+12B/\pi^2-2}{2\ln(\ln(x))} - \frac{\ln(3)-\ln(\ln(\ln(x)))}{2\ln(\ln(x))} \right)$$

where

$$B = - \int_0^\infty \ln(1 - e^{-y}) \ln(y) dy = \zeta'(2) - \frac{\pi^2}{6}\gamma.$$

**2.2. Meissel-Mertens Constants.** See [87] for more occurrences of the constants  $M$  and  $M'$ , and [88] for a historical treatment. Higher-order asymptotic series for  $E_n(\omega)$ ,  $\text{Var}_n(\omega)$ ,  $E_n(\Omega)$  and  $\text{Var}_n(\Omega)$  are given in [89]. While  $\sum_p 1/p$  is divergent, the following prime series is convergent [90]:

$$\sum_p \left( \frac{1}{p^2} + \frac{1}{p^3} + \frac{1}{p^4} + \dots \right) = \sum_p \frac{1}{p(p-1)} = 0.7731566690\dots$$

Also, the reciprocal sum of products of two primes satisfies [91]

$$\lim_{n \rightarrow \infty} \left( \sum_{pq \leq n} \frac{1}{pq} - \ln(\ln(n))^2 - 2M \ln(\ln(n)) \right) = \frac{\pi^2}{6} + M^2$$

and we wonder about the numerical values of

$$\begin{aligned} \sum_{p,q} \sum_{k=2}^\infty \frac{1}{(pq)^k} &= \sum_{p,q} \frac{1}{pq(pq-1)}, \\ \sum_{p,q} \sum_{k=2}^\infty \frac{1}{k(pq)^k} &= - \sum_{p,q} \left( \ln \left( 1 - \frac{1}{pq} \right) + \frac{1}{pq} \right). \end{aligned}$$

The second moment of  $\text{Im}(\ln(\zeta(1/2 + it)))$  over an interval  $[0, T]$  involves asymptotically a constant [92, 93]

$$\sum_{m=2}^\infty \sum_p \left( \frac{1}{m} - \frac{1}{m^2} \right) \frac{1}{p^m} = - \sum_p \left( \ln \left( 1 - \frac{1}{p} \right) + \text{Li}_2 \left( \frac{1}{p} \right) \right) = 0.1762478124\dots$$

as  $T \rightarrow \infty$ . If  $Q_k$  denotes the set of positive integers  $n$  for which  $\Omega(n) - \omega(n) = k$ , then  $Q_1 = \tilde{S}$  and the asymptotic density  $\delta_k$  satisfies [94, 95, 96]

$$\lim_{k \rightarrow \infty} 2^k \delta_k = \frac{1}{4C_{\text{twin}}} = 0.3786950320\dots;$$

the expression  $4C_{\text{twin}}$  also appears on pages 86 and 133–134, as well as the preceding entry.

Define  $f_k(n) = \#\{p : p^k | n\}$  and  $F_k(n) = \#\{p^{k+m} : p^{k+m} | n \text{ and } m \geq 0\}$ ; hence  $f_1(n) = \omega(n)$  and  $F_1(n) = \Omega(n)$ . It is known that, for  $k \geq 2$ ,

$$\sum_{n \leq x} f_k(n) \sim x \sum_p \frac{1}{p^k}, \quad \sum_{n \leq x} F_k(n) \sim x \sum_p \frac{1}{p^{k-1}(p-1)}$$

as  $x \rightarrow \infty$ . Also define  $g_k(n) = \#\{p : p | n \text{ and } p^k \nmid n\}$  and  $G_k(n) = \#\{p^m : p^m | n, p^k \nmid n \text{ and } m \geq 1\}$ . Then, for  $k \geq 2$ ,

$$\sum_{n \leq x} g_k(n) \sim x \left( \ln(\ln(x)) + M - \sum_p \frac{1}{p^k} \right),$$

$$\sum_{n \leq x} G_k(n) \sim x \left( \ln(\ln(x)) + M + \sum_p \frac{p^{k-1} - kp + k - 1}{p^k(p-1)} \right)$$

as  $x \rightarrow \infty$ . Other variations on  $k$ -full and  $k$ -free prime factors appear in [97]; the growth rate of  $\sum_{n \leq x} 1/\omega(n)$  and  $\sum_{n \leq x} 1/\Omega(n)$  is covered in [98] as well.

**2.3. Landau-Ramanujan Constant.** Define  $B_{3,j}(x)$  to be the number of positive integers  $\leq x$ , all of whose prime factors are  $\equiv j \pmod{3}$ , where  $j = 1$  or  $2$ . We have [99, 100, 101]

$$\lim_{x \rightarrow \infty} \frac{\sqrt{\ln(x)}}{x} B_{3,1}(x) = \frac{\sqrt{3}}{9K_3} = 0.3012165544\dots,$$

$$\lim_{x \rightarrow \infty} \frac{\sqrt{\ln(x)}}{x} B_{3,2}(x) = \frac{2\sqrt{3}K_3}{\pi} = 0.7044984335\dots$$

Here is a more complicated example (which arises in the theory of partitions). Let

$$W(x) = \#\left\{n \leq x : n = 2^h p_1^{e_1} p_2^{e_2} \cdots p_h^{e_h}, h \geq 1, e_k \geq 1, p_k \equiv 3, 5, 6 \pmod{7} \text{ for all } k\right\},$$

then the Selberg-Delange method gives [102, 103]

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{\ln(x)^{3/4}}{x} W(x) &= \frac{1}{\Gamma(1/4)} \left( \frac{6}{\sqrt{7}\pi} \right)^{1/4} \prod_{\substack{p \equiv 3, 5, 6 \\ \pmod{7}}} \left( 1 + \frac{1}{2(p-1)} \right) \left( 1 - \frac{1}{p} \right)^{1/4} \left( 1 + \frac{1}{p} \right)^{-1/4} \\ &= \frac{1}{\Gamma(1/4)} \left( \frac{6}{\sqrt{7}\pi} \right)^{1/4} (1.0751753443\dots) = 0.2733451113\dots \\ &= \frac{7}{24} (0.9371832387\dots). \end{aligned}$$

Other examples appear in [103] as well. We mention that products like [104]

$$\prod_{p \equiv 3 \pmod{4}} \left( 1 - \frac{2p}{(p^2 + 1)(p - 1)} \right) = 0.6436506796\dots,$$

$$\prod_{p \equiv 2 \pmod{3}} \left( 1 - \frac{2p}{(p^2 + 1)(p - 1)} \right) = 0.1739771224\dots$$

are evaluated to high precision in [105, 106] via special values of Dirichlet L-series.

**2.4. Artin's Constant.** Stephens' constant 0.5759... and Matthews' constant 0.1473... actually first appeared in [107]. Let  $\iota(n) = 1$  if  $n$  is square-free and  $\iota(n) = 0$  otherwise. Then [108, 109, 110, 111, 112, 113, 114]

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \iota(n)\iota(n+1) &= \prod_p \left( 1 - \frac{2}{p^2} \right) = 0.3226340989\dots = -1 + 2(0.6613170494\dots) \\ &= \frac{6}{\pi^2} \prod_p \left( 1 - \frac{1}{p^2 - 1} \right) = \frac{6}{\pi^2} (0.5307118205\dots), \end{aligned}$$

that is, the Feller-Tornier constant arises with regard to consecutive square-free numbers and to other problems. Also, consider the cardinality  $N(X)$  of nontrivial primitive integer vectors  $(x_0, x_1, x_2, x_3)$  that fall on Cayley's cubic surface

$$x_0x_1x_2 + x_0x_1x_3 + x_0x_2x_3 + x_1x_2x_3 = 0$$

and satisfy  $|x_j| \leq X$  for  $0 \leq j \leq 3$ . It is known that  $N(X) \sim cX(\ln(X))^6$  for some constant  $c > 0$  [115, 116]; finding  $c$  remains an open problem.

**2.5. Hafner-Sarnak-McCurley Constant.** In the "Added In Press" section (pages 601–602), the asymptotics of coprimality and of square-freeness are discussed for the Gaussian integers and for the Eisenstein-Jacobi integers. Cai & Bach [117] and Tóth [118] independently proved that the probability that  $k$  positive integers are *pairwise* coprime is [119]

$$\prod_p \left( 1 - \frac{1}{p} \right)^{k-1} \left( 1 + \frac{k-1}{p} \right) = \lim_{N \rightarrow \infty} \frac{(k-1)!}{N \ln(N)^{k-1}} \sum_{n=1}^N k^{\omega(n)}.$$

Freiberg [120], building on Moree's work [121], determined the probability that three positive integers are pairwise *not* coprime to be  $1 - 18/\pi^2 + 3P - Q = 0.1742197830\dots$ . More about sums involving  $2^{\omega(n)}$  and  $2^{-\omega(n)}$  appears in [122]. The asymptotics of  $\sum_{n=1}^N 3^{\Omega(n)}$ , due to Tenenbaum, are mentioned in [89]. Also, we have [123]

$$\sum_{n \leq N} \kappa(n)^\ell \sim \frac{1}{\ell + 1} \frac{\zeta(2\ell + 2)}{\zeta(2)} N^{\ell+1},$$



$$\sum_{n \leq N} K(n)^\ell \sim \frac{1}{\ell+1} \frac{\zeta(\ell+1)}{\zeta(2)} \prod_p \left(1 - \frac{1}{p^\ell(p+1)}\right) \cdot N^{\ell+1}$$

as  $N \rightarrow \infty$ , for any positive integer  $\ell$ . In the latter formula, the product for  $\ell = 1$  and  $\ell = 2$  appears in [122] with regard to the number/sum of unitary square-free divisors; the product for  $\ell = 2$  further is connected with class number theory [69].

**2.6. Niven's Constant.** The quantity  $C$  appears unexpectedly in [124]. If we instead examine the mean of the exponents:

$$L(m) = \begin{cases} 1 & \text{if } m = 1, \\ \frac{1}{k} \sum_{j=1}^k a_j & \text{if } m > 1, \end{cases}$$

then [125, 126]

$$\sum_{m \leq n} L(m) = n + C_1 \frac{n}{\ln(\ln(n))} + C_2 \frac{n}{\ln(\ln(n))^2} + O\left(\frac{n}{\ln(\ln(n))^3}\right)$$

as  $n \rightarrow \infty$ , where [90]

$$C_1 = \sum_p \frac{1}{p(p-1)} = M' - M = 0.7731566690\dots,$$

$$C_2 = \sum_p \frac{1}{p^2(p-1)} - C_1 M = C_1(1 - M) - N = 0.1187309349\dots,$$

using notation defined on pages 94–95. The constant  $C_1$  also appears in our earlier entry [2.2]. We will report on [127] soon.

**2.7. Euler Totient Constants.** Let us clarify the third sentence:  $\varphi(n)$  is the number of generators in  $\mathbb{Z}_n$ , the additive group of integers modulo  $n$ . It is also the number of elements in  $\mathbb{Z}_n^*$ , the multiplicative group of invertible integers modulo  $n$ .

Define  $f(n) = n\varphi(n)^{-1} - e^\gamma \ln(\ln(n))$ . Nicolas [128] proved that  $f(n) > 0$  for infinitely many integers  $n$  by the following reasoning. Let  $P_k$  denote the product of the first  $k$  prime numbers. If the Riemann hypothesis is true, then  $f(P_k) > 0$  for all  $k$ . If the Riemann hypothesis is false, then  $f(P_k) > 0$  for infinitely many  $k$  and  $f(P_l) \leq 0$  for infinitely many  $l$ .

Let  $U(n)$  denote the set of values  $\leq n$  taken by  $\varphi$  and  $v(n)$  denote its cardinality; for example [129],  $U(15) = \{1, 2, 4, 6, 8, 10, 12\}$  and  $v(15) = 7$ . Let  $\ln_2(x) = \ln(\ln(x))$  and  $\ln_m(x) = \ln(\ln_{m-1}(x))$  for convenience. Ford [130] proved that

$$v(n) = \frac{n}{\ln(n)} \exp \left\{ C[\ln_3(n) - \ln_4(n)]^2 + D \ln_3(n) - \left[ D + \frac{1}{2} - 2C \right] \ln_4(n) + O(1) \right\}$$

as  $n \rightarrow \infty$ , where

$$C = -\frac{1}{2\ln(\rho)} = 0.8178146464\dots,$$

$$D = 2C(1 + \ln(F'(\rho)) - \ln(2C)) - \frac{3}{2} = 2.1769687435\dots$$

$$F(x) = \sum_{k=1}^{\infty} ((k+1)\ln(k+1) - k\ln(k) - 1)x^k$$

and  $\rho = 0.5425985860\dots$  is the unique solution on  $[0, 1)$  of the equation  $F(\rho) = 1$ . Also,

$$\lim_{n \rightarrow \infty} \frac{1}{v(n)\ln_2(n)} \sum_{m \in U(n)} \omega(m) = \frac{1}{1-\rho} = 2.1862634648\dots$$

which contrasts with a related result of Erdős & Pomerance [131]:

$$\lim_{n \rightarrow \infty} \frac{1}{n \ln_2(n)^2} \sum_{m=1}^n \omega(\varphi(n)) = \frac{1}{2}.$$

These two latter formulas hold as well if  $\omega$  is replaced by  $\Omega$ . See [132] for more on Euler's totient.

Define the *reduced totient* or *Carmichael function*  $\psi(n)$  to be the size of the largest cyclic subgroup of  $\mathbb{Z}_n^*$ . We have [133]

$$\frac{1}{N} \sum_{n \leq N} \psi(n) = \frac{N}{\ln(N)} \exp \left[ \frac{P \ln_2(N)}{\ln_3(N)} (1 + o(1)) \right]$$

as  $N \rightarrow \infty$ , where

$$P = e^{-\gamma} \prod_p \left( 1 - \frac{1}{(p-1)^2(p+1)} \right) = 0.3453720641\dots$$

There is a set  $S$  of positive integers of asymptotic density 1 such that, for  $n \in S$ ,

$$n\psi(n)^{-1} = (\ln(n))^{\ln_3(n)+Q+o(1)}$$

and

$$Q = -1 + \sum_p \frac{\ln(p)}{(p-1)^2} = 0.2269688056\dots;$$

it is not known whether  $S = \mathbb{Z}^+$  is possible.

**2.8. Pell-Steinhagen Constants.** The constant  $P$  is transcendental via a general theorem on values of modular forms due to Nesterenko [134, 135]. Here is a

constant similar to  $P$ : The number of positive integers  $n \leq N$ , for which  $2n - 1$  is not divisible by  $2^p - 1$  for any prime  $p$ , is  $\sim cN$ , where

$$c = \prod_p \left(1 - \frac{1}{2^p - 1}\right) = 0.5483008312\dots$$

A ring-theoretic analog of this statement, plus generalizations, appear in [136].

**2.10. Sierpinski's Constant.** Sierpinski's formulas for  $\hat{S}$  and  $\tilde{S}$  contained a few errors: they should be [137, 138, 139, 140, 141, 142]

$$\hat{S} = \gamma + S - \frac{12}{\pi^2}\zeta'(2) + \frac{\ln(2)}{3} - 1 = 1.7710119609\dots = \frac{\pi}{4}(2.2549224628\dots),$$

$$\tilde{S} = 2S - \frac{12}{\pi^2}\zeta'(2) + \frac{\ln(2)}{3} - 1 = 2.0166215457\dots = \frac{1}{4}(8.0664861829\dots).$$

In the summation formula at the top of page 125,  $D_n$  should be  $D_k$ . Also, the divisor analog of Sierpinski's second series is [143]

$$\sum_{k=1}^n d(k^2) = \left(\frac{3}{\pi^2} \ln(n)^2 + \left(\frac{18\gamma - 6}{\pi^2} - \frac{72}{\pi^4}\zeta'(2)\right) \ln(n) + c\right) n + O(n^{1/2+\varepsilon})$$

as  $n \rightarrow \infty$ , where the expression for  $c$  is complicated. It is easily shown that  $d(n^2)$  is the number of ordered pairs of positive integers  $(i, j)$  satisfying  $\text{lcm}(i, j) = n$ .

Define  $R(n)$  to be the number of representations of  $n$  as a sum of three squares, counting order and sign. Then

$$\sum_{k=1}^n R(k) = \frac{4\pi}{3}n^{3/2} + O(n^{3/4+\varepsilon})$$

for all  $\varepsilon > 0$  and [144]

$$\sum_{k=1}^n R(k)^2 = \frac{8\pi^4}{21\zeta(3)}n^2 + O(n^{14/9}).$$

The former is the same as the number of integer ordered triples falling within the ball of radius  $\sqrt{n}$  centered at the origin; an extension of the latter to sums of  $m$  squares, when  $m > 3$ , is also known [144].

**2.11. Abundant Numbers Density Constant.** The definition of  $A(x)$  should be replaced by

$$A(x) = \lim_{n \rightarrow \infty} \frac{|\{k \leq n : \sigma(k) \geq xk\}|}{n}.$$

If  $K(x)$  is the number of all positive integers  $m$  that satisfy  $\sigma(m) \leq x$ , then [145]

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{K(x)}{x} &= \prod_p \left(1 - \frac{1}{p}\right) \left(1 + \sum_{j=1}^{\infty} \left(1 + \sum_{i=1}^j p^i\right)^{-1}\right) \\ &= \prod_p \left(1 - \frac{1}{p}\right) \left(1 + (p-1) \sum_{j=1}^{\infty} \frac{1}{p^{j+1} - 1}\right). \end{aligned}$$

**2.12. Linnik's Constant.** In the definition of  $L$ , “lim” should be replaced by “limsup”. Clearly  $L$  exists; the fact that  $L < \infty$  was Linnik's important contribution.

**2.13. Mills' Constant.** Let  $q_1 < q_2 < \dots < q_k$  denote the consecutive prime factors of an integer  $n > 1$ . Define

$$F(n) = \sum_{j=1}^{k-1} \left(1 - \frac{q_j}{q_{j+1}}\right) = \omega(n) - 1 - \sum_{j=1}^{k-1} \frac{q_j}{q_{j+1}}$$

if  $k > 1$  and  $F(n) = 0$  if  $k = 1$ . Erdős & Nicolas [146] demonstrated that there exists a constant  $C' = 1.70654185\dots$  such that, as  $n \rightarrow \infty$ ,  $F(n) \leq \sqrt{\ln(n)} - C' + o(1)$ , with equality holding for infinitely many  $n$ . Further,  $C' = C + \ln(2) + 1/2$ , where [146, 147]

$$C = \sum_{i=1}^{\infty} \left\{ \ln \left( \frac{p_{i+1}}{p_i} \right) - \left( 1 - \frac{p_i}{p_{i+1}} \right) \right\} = 0.51339467\dots, \quad \sum_{i=1}^{\infty} \left( \frac{p_{i+1}}{p_i} - 1 \right)^2 = 1.65310351\dots,$$

and  $p_1 = 2, p_2 = 3, p_3 = 5, \dots$  is the sequence of all primes.

It now seems that  $\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) / \ln(p_n) = 0$  is a theorem [148, 149], clarifying the uncertainty raised in “Added In Press” (pages 601–602). More about small prime gaps will surely appear soon; research concerning large prime gaps continues as well [150, 151].

**2.15. Glaisher-Kinkelin Constant.** In the second display for  $D(x)$ ,  $\exp(-x/2)$  should be replaced by  $\exp(x/2)$ . Another proof of the formula for  $D(1)$  is given in [47]; another special case is [152]

$$D(1/2) = \frac{2^{1/6} \sqrt{\pi} A^3}{\Gamma(1/4)} e^{G/\pi}.$$

The two quantities

$$G_2 \left( \frac{1}{2} \right) = 0.6032442812\dots, \quad G_2 \left( \frac{3}{2} \right) = \sqrt{\pi} G_2 \left( \frac{1}{2} \right) = 1.0692226492\dots$$

play a role in a discussion of the limiting behavior of Toeplitz determinants and the Fisher-Hartwig conjecture [153, 154]. Kraovskiy [155] and Ehrhardt [156] proved

Dyson's conjecture regarding the asymptotic expansion of  $E(s)$  as  $s \rightarrow \infty$ ; a third proof is given in [157]. Also, the quantities

$$G_2 \left(\frac{1}{2}\right)^{-1} = 1.6577032408\dots = 2^{-1/24} e^{-3/16} \pi^{1/4} (3.1953114860\dots)^{3/8}$$

$$G_3 \left(\frac{3}{2}\right)^{-1} = G_2 \left(\frac{1}{2}\right) G_3 \left(\frac{1}{2}\right)^{-1} = 0.9560900097\dots = \pi^{-1/2} (3.3388512141\dots)^{7/16}$$

appear in [158]. In the last paragraph on page 141, the polynomial  $q(x)$  should be assumed to have degree  $n$ . See [159, 160] for more on the GUE hypothesis.

**2.16. Stolarsky-Harboth Constant.** Given a positive integer  $n$ , define  $s_1^2$  to be the largest square not exceeding  $n$ . Then define  $s_2^2$  to be the largest square not exceeding  $n - s_1^2$ , and so forth. Hence  $n = \sum_{j=1}^r s_j^2$  for some  $r$ . We say that  $n$  is a *greedy sum of distinct squares* if  $s_1 > s_2 > \dots > s_r$ . Let  $A(N)$  be the number of such integers  $n < N$ , plus one. Montgomery & Vorhauer [161] proved that  $A(N)/N$  does not tend to a constant, but instead that there is a continuous function  $f(x)$  of period 1 for which

$$\lim_{k \rightarrow \infty} \frac{A(4 \exp(2^{k+x}))}{4 \exp(2^{k+x})} = f(x), \quad \min_{0 \leq x \leq 1} f(x) = 0.50307\dots < \max_{0 \leq x \leq 1} f(x) = 0.50964\dots$$

where  $k$  takes on only integer values. This is reminiscent of the behavior discussed for digital sums. Two simple examples, due to Hardy [162, 163] and Elkies [164], involve the series

$$\varphi(x) = \sum_{k=0}^{\infty} x^{2^k}, \quad \psi(x) = \sum_{k=0}^{\infty} (-1)^k x^{2^k}.$$

As  $x \rightarrow 1^-$ , the asymptotics of  $\varphi(x)$  and  $\psi(x)$  are complicated by oscillating errors with amplitude

$$\sup_{x \rightarrow 1^-} \left| \varphi(x) + \frac{\ln(-\ln(x)) + \gamma}{\ln(2)} - \frac{3}{2} + x \right| = (1.57\dots) \times 10^{-6},$$

$$\sup_{x \rightarrow 1^-} \left| \psi(x) - \frac{1}{6} - \frac{1}{3}x \right| = (2.75\dots) \times 10^{-3}.$$

The function  $\varphi(x)$  also appears in what is known as Catalan's integral (section 1.5.2) for Euler's constant  $\gamma$ . See [165] as well.

**2.17. Gauss-Kuzmin-Wirsing Constant.** The preprint math.NT/9908043 was withdrawn by the author without comment; the bounds we gave for the Hausdorff dimension of real numbers with partial denominators in  $\{1, 2\}$  remain unaffected.

**2.18. Porter-Hensley Constants.** Lhote [166, 167] developed rigorous techniques for computing certain variances to high precision, for example,  $4\lambda_1''(2)$ . With

regard to the binary GCD algorithm, Maze [168] confirmed Brent's functional equation for a certain limiting distribution [169]

$$g(x) = \sum_{k \geq 1} 2^{-k} \left( g \left( \frac{1}{1 + 2^k/x} \right) - g \left( \frac{1}{1 + 2^k x} \right) \right), \quad 0 \leq x \leq 1$$

as well as the formula

$$2 + \frac{1}{\ln(2)} \int_0^1 \frac{g(x)}{1-x} dx = \frac{2}{\kappa \ln(2)} = 2.8329765709\dots = \frac{\pi^2(0.3979226811\dots)}{2 \ln(2)}.$$

**2.20. Erdős' Reciprocal Sum Constants.** A sequence of positive integers  $b_1 < b_2 < \dots < b_m$  is a  $B_h$ -sequence if all  $h$ -fold sums  $b_{i_1} + b_{i_2} + \dots + b_{i_h}$ ,  $i_1 \leq i_2 \leq \dots \leq i_h$ , are distinct. Given  $n$ , choose a  $B_h$ -sequence  $\{b_i\}$  so that  $b_m \leq n$  and  $m$  is maximal; let  $F_h(n)$  be this value of  $m$ . It is known that  $C_h = \limsup_{n \rightarrow \infty} n^{-1/h} F_h(n)$  is finite; we further have [170, 171, 172, 173, 174, 175]

$$C_2 = 1, \quad 1 \leq C_3 \leq (7/2)^{1/3}, \quad 1 \leq C_4 \leq 7^{1/4}.$$

More generally, a sequence of positive integers  $b_1 < b_2 < \dots < b_m$  is a  $B_{h,g}$ -sequence if, for every positive integer  $k$ , the equation  $x_1 + x_2 + \dots + x_h = k$ ,  $x_1 \leq x_2 \leq \dots \leq x_h$ , has at most  $g$  solutions with  $x_j = b_{i_j}$  for all  $j$ . Defining  $F_{h,g}(n)$  and  $C_{h,g}$  analogously, we have [175, 176, 177, 178, 179]

$$\frac{4\sqrt{7}}{7} \leq C_{2,2} \leq \frac{\sqrt{21}}{2}, \quad \frac{3\sqrt{2}}{4} g^{1/2} + o(g^{1/2}) \leq C_{2,g} \leq \min \left\{ \frac{7}{2} g - \frac{7}{4}, \frac{17g}{5} \right\}^{1/2}$$

as  $g \rightarrow \infty$ .

**2.21. Stieltjes Constants.** If  $d_k(n)$  denotes the number of sequences  $x_1, x_2, \dots, x_k$  of positive integers such that  $n = x_1 x_2 \dots x_k$ , then [180, 181, 182]

$$\sum_{n=1}^N d_2(n) \sim N \ln(N) + (2\gamma_0 - 1)N \quad (d_2 \text{ is the divisor function}),$$

$$\sum_{n=1}^N d_3(n) \sim \frac{1}{2} N \ln(N)^2 + (3\gamma_0 - 1)N \ln(N) + (-3\gamma_1 + 3\gamma_0^2 - 3\gamma_0 + 1)N,$$

$$\begin{aligned} \sum_{n=1}^N d_4(n) &\sim \frac{1}{6} N \ln(N)^3 + \frac{4\gamma_0 - 1}{2} N \ln(N)^2 + (-4\gamma_1 + 6\gamma_0^2 - 4\gamma_0 + 1)N \ln(N) \\ &\quad + (2\gamma_2 - 12\gamma_1\gamma_0 + 4\gamma_1 + 4\gamma_0^3 - 6\gamma_0^2 + 4\gamma_0 - 1)N \end{aligned}$$

as  $N \rightarrow \infty$ . More generally,  $\sum_{n=1}^N d_k(n)$  can be asymptotically expressed as  $N$  times a polynomial of degree  $k - 1$  in  $\ln(N)$ , which in turn can be described as the residue at  $z = 1$  of  $z^{-1}\zeta(z)^k N^z$ . See [89] for an application of  $\{\gamma_j\}_{j=0}^{\infty}$  to asymptotic series for  $E_n(\omega)$  and  $E_n(\Omega)$ , and [183, 184, 185, 186, 187, 188] for connections to the Riemann hypothesis.

**2.23. Diophantine Approximation Constants.** Which planar, symmetric, bounded convex set  $K$  has the worst packing density? If  $K$  is a disk, the packing density is  $\pi/\sqrt{12} = 0.9068996821\dots$ , which surprisingly is better than if  $K$  is the smoothed octagon:

$$\frac{8 - 4\sqrt{2} - \ln(2)}{2\sqrt{2} - 1} = \frac{1}{4}(3.6096567319\dots) = 0.9024141829\dots$$

Do worse examples exist? The answer is only conjectured to be yes [189].

**2.25. Cameron's Sum-Free Set Constants.** Erdős [190] and Alon & Kleitman [191] showed that any finite set  $B$  of positive integers must contain a sum-free subset  $A$  such that  $|A| > \frac{1}{3}|B|$ . See also [192, 193, 194]. The largest constant  $c$  such that  $|A| > c|B|$  must satisfy  $1/3 \leq c < 12/29$ , but its exact value is unknown. Using harmonic analysis, Bourgain [195] improved the original inequality to  $|A| > \frac{1}{3}(|B|+2)$ . Green [196, 197] demonstrated that  $s_n = O(2^{n/2})$ , but the values  $c_o = 6.8\dots$  and  $c_e = 6.0\dots$  await more precise computation.

Further evidence for the existence of complete aperiodic sum-free sets is given in [198].

**2.29. Fast Matrix Multiplication Constants.** Efforts continue [199, 200] to reduce the upper bound on  $\omega$  to 2.

**2.30. Pisot-Vijayaraghavan-Salem Constants.** The definition of Mahler's measure  $M(\alpha)$  is unclear: It should be the product of  $\max\{1, |\alpha_j|\}$  over all  $1 \leq j \leq n$ . Breusch [201] gave a lower bound  $> 1$  for  $M(\alpha)$  of non-reciprocal algebraic integers  $\alpha$ , anticipating Smyth's stronger result by twenty years.

Compare the sequence  $\{(3/2)^n\}$ , for which little is known, with the recursion  $x_0 = 0$ ,  $x_n = \{x_{n-1} + \ln(3/2)/\ln(2)\}$ , for which a musical interpretation exists. If a guitar player touches a vibrating string at a point two-thirds from the end of the string, its fundamental frequency is dampened and a higher overtone is heard instead. This new pitch is a perfect fifth above the original note. It is well-known that the "circle of fifths" never closes, in the sense that  $2^{x_n}$  is never an integer for  $n > 0$ . Further, the "circle of fifths", in the limit as  $n \rightarrow \infty$ , fills the continuum of pitches spanning the octave [202, 203].

The Collatz function  $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$  is defined by

$$f(n) = \begin{cases} 3n + 1 & \text{if } n \text{ is odd} \\ n/2 & \text{if } n \text{ is even} \end{cases} .$$

Let  $f^k$  denote the  $k^{\text{th}}$  iterate of  $f$ . The  $3x + 1$  conjecture asserts that, given any positive integer  $n$ , there exists  $k$  such that  $f^k(n) = 1$ . Let  $\sigma(n)$  be the first  $k$  such that  $f^k(n) < n$ , called the *stopping time* of  $n$ . If we could demonstrate that every positive integer  $n$  has a finite stopping time, then the  $3x + 1$  conjecture would be proved. Heuristic reasoning [204, 205, 206] provides that the average stopping time over all odd integers  $1 \leq n \leq N$  is asymptotically

$$\lim_{N \rightarrow \infty} E_{\text{odd}}(\sigma(n)) = \sum_{j=1}^{\infty} \left[ 1 + \left( 1 + \frac{\ln(3)}{\ln(2)} \right) j \right] c_j 2^{-\lfloor \frac{\ln(3)}{\ln(2)} j \rfloor} = 9.4779555565\dots$$

where  $c_j$  is the number of admissible sequences of order  $j$ . Such a sequence  $\{a_k\}_{k=1}^m$  satisfies  $a_k = 3/2$  exactly  $j$  times,  $a_k = 1/2$  exactly  $m - j$  times,  $\prod_{k=1}^m a_k < 1$  but  $\prod_{k=1}^l a_k > 1$  for all  $1 \leq l < m$  [207]. In contrast, the *total* stopping time  $\sigma_{\infty}(n)$  of  $n$ , the first  $k$  such that  $f^k(n) = 1$ , appears to obey

$$\lim_{N \rightarrow \infty} E \left( \frac{\sigma_{\infty}(n)}{\ln(n)} \right) \sim \frac{2}{2 \ln(2) - \ln(3)} = 6.9521189935\dots = \frac{2}{\ln(10)} (8.0039227796\dots).$$

**2.32. De Bruijn-Newman Constant.** Further work regarding Li's criterion, which is equivalent to Riemann's hypothesis and which involves the Stieltjes constants, appears in [183, 184]. A different criterion is due to Matiyasevich [185, 186]; the constant  $-\ln(4\pi) + \gamma + 2 = 0.0461914179\dots = 2(0.0230957089\dots)$  comes out as a special case. See also [187, 188]. As another aside, we mention the unboundedness of  $\zeta(1/2 + it)$  for  $t \in (0, \infty)$ , but that a precise order of growth remains open [208, 209, 210]. In contrast, there is a conjecture that [211, 212]

$$\max_{t \in [T, 2T]} |\zeta(1 + it)| = e^{\gamma} (\ln(\ln(T)) + \ln(\ln(\ln(T))) + C + o(1)),$$

$$\max_{t \in [T, 2T]} \frac{1}{|\zeta(1 + it)|} = \frac{6e^{\gamma}}{\pi^2} (\ln(\ln(T)) + \ln(\ln(\ln(T))) + C + o(1))$$

as  $T \rightarrow \infty$ , where

$$C = 1 - \ln(2) + \int_0^2 \frac{\ln(I_0(t))}{t^2} dt + \int_2^{\infty} \frac{\ln(I_0(t)) - t}{t^2} dt = -0.0893\dots$$

and  $I_0(t)$  is the zeroth modified Bessel function. These formulas have implications for  $|\zeta(it)|$  and  $1/|\zeta(it)|$  as well by the analytic continuation formula.

**2.33. Hall-Montgomery Constant.** Let  $\psi$  be the unique solution on  $(0, \pi)$  of the equation  $\sin(\psi) - \psi \cos(\psi) = \pi/2$  and define  $K = -\cos(\psi) = 0.3286741629\dots$



Consider any real multiplicative function  $f$  whose values are constrained to  $[-1, 1]$ . Hall & Tenenbaum [213] proved that, for some constant  $C > 0$ ,

$$\sum_{n=1}^N f(n) \leq CN \exp \left\{ -K \sum_{p \leq N} \frac{1 - f(p)}{p} \right\} \quad \text{for sufficiently large } N,$$

and that, moreover, the constant  $K$  is sharp. (The latter summation is over all prime numbers  $p$ .) This interesting result is a lemma used in [214]. A table of values of sharp constants  $K$  is also given in [213] for the generalized scenario where  $f$  is complex,  $|f| \leq 1$  and, for all primes  $p$ ,  $f(p)$  is constrained to certain elliptical regions in  $\mathbb{C}$ .

**3.3. Landau-Kolmogorov Constants.** For  $L_2(0, \infty)$ , Bradley & Everitt [215] were the first to determine that  $C(4, 2) = 2.9796339059\dots = \sqrt{8.8782182137\dots}$ ; see also [216, 217, 218]. Ditzian [219] proved that the constants for  $L_1(-\infty, \infty)$  are the same as those for  $L_\infty(-\infty, \infty)$ . Phóng [217] obtained the following best possible inequality in  $L_2(0, 1)$ :

$$\int_0^1 |f'(x)|^2 dx \leq (6.4595240299\dots) \left( \int_0^1 |f(x)|^2 dx + \int_0^1 |f''(x)|^2 dx \right)$$

where the constant is given by  $\sec(2\theta)/2$  and  $\theta$  is the unique zero satisfying  $0 < \theta < \pi/4$  of

$$\begin{aligned} & \sin(\theta)^4 \left( e^{2\sin(\theta)} - 1 \right)^2 \left( e^{-2\sin(\theta)} - 1 \right)^2 + \cos(\theta)^4 [2 - 2\cos(2\cos(\theta))]^2 \\ & - \cos(2\theta)^4 [1 + e^{4\sin(\theta)} - 2e^{2\sin(\theta)} \cos(2\cos(\theta))] [1 + e^{-4\sin(\theta)} - 2e^{-2\sin(\theta)} \cos(2\cos(\theta))] \\ & - 2\cos(\theta)^2 \sin(\theta)^2 [2 - 2\cos(2\cos(\theta))] (1 - e^{-2\sin(\theta)}) \left( e^{2\sin(\theta)} - 1 \right). \end{aligned}$$

We wonder about other such additive analogs of Landau-Kolmogorov inequalities.

**3.5. Copson-de Bruijn Constant.** More relevant material is found in Ackermans [220].

**3.6. Sobolev Isoperimetric Constants.** In section 3.6.1,  $\sqrt{\lambda} = 1$  represents the principal frequency of the sound we hear when a string is plucked; in section 3.6.3,  $\sqrt{\lambda} = \theta$  represents likewise when a kettledrum is struck. (The square root was missing in both.) The units of frequency, however, are not compatible between these two examples.

The “rod” constant  $500.5639017404\dots = (4.7300407448\dots)^4$  appears in [221, 222, 223]. It is the second term in a sequence  $c_1, c_2, c_3, \dots$  for which  $c_1 = \pi^2 = 9.869\dots$  (in connection with the “string” inequality) and  $c_3 = (2\pi)^6 = 61528.908\dots$ ; the constant  $c_4$  is the smallest eigenvalue of ODE

$$f^{(viii)}(x) = \lambda f(x), \quad 0 \leq x \leq 1,$$

$$f(0) = f'(0) = f''(0) = f'''(0) = 0, \quad f(1) = f'(1) = f''(1) = f'''(1) = 0$$

and evidently is not known. Allied subjects include positive definite Toeplitz matrices and conditioning of certain least squares problems.

More relevant material is found in [224, 225, 226].

**3.10. Kneser-Mahler Constants.** The constants  $\ln(\beta)$  and  $\ln(\delta)$  appear in [227]. Conjectured L-series expressions for  $M\left(1 + \sum_{j=1}^n x_j\right)$ , due to Rodriguez-Villegas, are exhibited for  $n = 4, 5$  in [228].

**3.12. Du Bois Reymond's Constants.** The constant  $(\pi/\xi)^2$  is equal to the largest eigenvalue of the infinite symmetric matrix  $(a_{m,n})_{m \geq 1, n \geq 1}$  with elements  $a_{m,n} = m^{-1}n^{-1} + m^{-2}\delta_{m,n}$ , where  $\delta_{m,n} = 1$  if  $m = n$  and  $\delta_{m,n} = 0$ . Boersma [229] employed this fact to give an alternative proof of Szegő's theorem.

**3.15. Van der Corput's Constant.** We examined only the case in which  $f$  is a real twice-continuously differentiable function on the interval  $[a, b]$ ; a generalization to the case where  $f$  is  $n$  times differentiable,  $n \geq 2$ , is discussed in [230, 231] with some experimental numerical results for  $n = 3$ .

**3.16. Turán's Power Sum Constants.** Recent work appears in [232, 233, 234, 235, 236, 237, 238, 239], to be reported on later.

**4.3. Achieser-Krein-Favard Constants.** While on the subject of trigonometric polynomials, we mention Littlewood's conjecture [240]. Let  $n_1 < n_2 < \dots < n_k$  be integers and let  $c_j$ ,  $1 \leq j \leq k$ , be complex numbers with  $|c_j| \geq 1$ . Konyagin [241] and McGehee, Pigno & Smith [242] proved that there exists  $C > 0$  so that the inequality

$$\int_0^1 \left| \sum_{j=1}^k c_j e^{2\pi i n_j \xi} \right| d\xi \geq C \ln(k)$$

always holds. It is known that the smallest such constant  $C$  satisfies  $C \leq 4/\pi^2$ ; Stegeman [243] demonstrated that  $C \geq 0.1293$  and Yabuta [244] improved this slightly to  $C \geq 0.129590$ . What is the true value of  $C$ ?

**4.5. The "One-Ninth" Constant.** Zudilin [245] deduced that  $\Lambda$  is transcendental by use of Theorem 4 in [246]. See also [247, 248].

**4.7. Berry-Esseen Constant.** Significant progress on the asymptotic case (as  $\lambda \rightarrow 0$ ) is described in [249, 250, 251]. A different form of the inequality is found in [252].

**4.8. Laplace Limit Constant.** The quantity  $\lambda = 0.6627434193\dots$  appears in [253] with regard to Plateau's problem for two circular rings dipped in soap solution. We hope to report on [254] at a later time.

**5.1. Abelian Group Enumeration Constants.** For a finite abelian group  $G$ , let  $r(G)$  denote the minimum number of generators of  $G$  and let  $E(G)$  denote the expected number of random elements from  $G$ , drawn independently and uniformly,

to generate  $G$ . Define  $e(G) = E(G) - r(G)$ , the *excess* of  $G$ . Then [124]

$$e_r = \sup \{e(G) : r(G) = r\} = 1 + \sum_{j=1}^{\infty} \left(1 - \prod_{k=1}^r \zeta(j+k)^{-1}\right);$$

in particular,  $e_1 = 1.7052111401\dots$  (Niven's constant) for the cyclic case and

$$\sigma = \lim_{r \rightarrow \infty} e_r = 1 + \sum_{j=2}^{\infty} \left(1 - \prod_{k=j}^{\infty} \zeta(k)^{-1}\right) = 2.118456563\dots$$

in general. It is remarkable that this limit is finite! Let also

$$\tau = \sum_{j=1}^{\infty} \left(1 - (1 - 2^{-j}) \prod_{k=j+1}^{\infty} \zeta(k)^{-1}\right) = 1.742652311\dots,$$

then for the multiplicative group  $\mathbb{Z}_n^*$  of integers relatively prime to  $n$ ,

$$\sup \{e(G) : G = \mathbb{Z}_n^* \text{ and } 2 < n \equiv l \pmod{8}\} = \begin{cases} \sigma & \text{if } l = 1, 3, 5 \text{ or } 7, \\ \sigma - 1 & \text{if } l = 2 \text{ or } 6, \\ \tau & \text{if } l = 4, \\ \tau + 1 & \text{if } l = 0. \end{cases}$$

We emphasize that  $l$ , not  $n$ , is fixed in the supremum (as according to the right-hand side). The constant  $A_1^{-1} = 0.4357570767\dots$  also appears as a ‘‘best probability’’ corresponding to certain nonabelian groups.

**5.4. Golomb-Dickman Constant.** Let  $P^+(n)$  denote the largest prime factor of  $n$  and  $P^-(n)$  denote the smallest prime factor of  $n$ . We mentioned that

$$\sum_{n=2}^N \ln(P^+(n)) \sim \lambda N \ln(N) - \lambda(1 - \gamma)N, \quad \sum_{n=2}^N \ln(P^-(n)) \sim e^{-\gamma} N \ln(\ln(N)) + cN$$

as  $N \rightarrow \infty$ , but did not give an expression for the constant  $c$ . Tenenbaum [255] found that

$$c = e^{-\gamma}(1 + \gamma) + \int_1^{\infty} \frac{\omega(t) - e^{-\gamma}}{t} dt + \sum_p \left\{ e^{-\gamma} \ln \left(1 - \frac{1}{p}\right) + \frac{\ln(p)}{p-1} \prod_{q \leq p} \left(1 - \frac{1}{q}\right) \right\},$$

where the sum over  $p$  and product over  $q$  are restricted to primes. A numerical evaluation is still open.

The longest tail  $L(\varphi)$ , given a random mapping  $\varphi : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ , is called the *height* of  $\varphi$  in [256, 257, 258] and satisfies

$$\lim_{n \rightarrow \infty} \mathbb{P} \left( \frac{L(\varphi)}{\sqrt{n}} \leq x \right) = \sum_{k=-\infty}^{\infty} (-1)^k \exp \left( -\frac{k^2 x^2}{2} \right)$$

for fixed  $x > 0$ . For example,

$$\lim_{n \rightarrow \infty} \text{Var} \left( \frac{L(\varphi)}{\sqrt{n}} \right) = \frac{\pi^2}{3} - 2\pi \ln(2)^2.$$

The longest rho-path  $R(\varphi)$  is called the *diameter* of  $\varphi$  in [259] and has moments

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \left( \frac{R(\varphi)}{\sqrt{n}} \right)^j \right] = \frac{\sqrt{\pi} j}{2^{j/2} \Gamma((j+1)/2)} \int_0^\infty x^{j-1} (1 - e^{\text{Ei}(-x) - I(x)}) dx$$

for fixed  $j > 0$ . Complicated formulas for the distribution of the largest tree  $P(\varphi)$  also exist [257, 258, 260].

A permutation  $p \in S_n$  is a *square* if  $p = q^2$  for some  $q \in S_n$ ; it is a *cube* if  $p = r^3$  for some  $r \in S_n$ . For convenience, let  $\omega = (-1 + i\sqrt{3})/2$  and

$$\Psi(x) = \frac{1}{3} \left( \exp(x) + 2 \exp(-x/2) \cos(\sqrt{3}x/2) \right).$$

The probability that a random  $n$ -permutation is a square is [261, 262, 263, 264, 265]

$$\begin{aligned} &\sim \frac{2^{1/2}}{\Gamma(1/2)} \frac{1}{n^{1/2}} \prod_{1 \leq m \equiv 0 \pmod{2}} \frac{e^{1/m} + e^{-1/m}}{2} = \sqrt{\frac{2}{\pi n}} \prod_{k=1}^{\infty} \cosh\left(\frac{1}{2k}\right) \\ &= \sqrt{\frac{2}{\pi n}} (1.2217795151\dots) = (0.9748390118\dots) n^{-1/2} \end{aligned}$$

as  $n \rightarrow \infty$ ; the probability that it is a cube is [264, 265]

$$\begin{aligned} &\sim \frac{3^{1/3}}{\Gamma(2/3)} \frac{1}{n^{1/3}} \prod_{1 \leq m \equiv 0 \pmod{3}} \frac{e^{1/m} + e^{\omega/m} + e^{\omega^2/m}}{3} \\ &= \frac{3^{5/6} \Gamma(1/3)}{2\pi n^{1/3}} \prod_{k=1}^{\infty} \Psi\left(\frac{1}{3k}\right) = (1.0729979443\dots) n^{-1/3}. \end{aligned}$$

Two permutations  $p, q \in S_n$  are of the *same cycle type* if their cycle decompositions are identical (in the sense that they possess the same number of cycles of length  $l$ , for each  $l \geq 1$ ). The probability that two independent, random  $n$ -permutations have the same cycle type is [265]

$$\sim \frac{1}{n^2} \prod_{k=1}^{\infty} I_0\left(\frac{2}{k}\right) = (4.2634035141\dots) n^{-2}$$

as  $n \rightarrow \infty$ , where  $I_0$  is the zeroth modified Bessel function.

More on the Erdős-Turán constant appears in [266, 267].

**5.5. Kalmár's Composition Constant.** See [268] for precise inequalities involving  $m(n)$  and  $\rho = 1.7286472389\dots$  A *Carlitz composition* of size  $n$  is an additive composition  $n = x_1 + x_2 + \dots + x_k$  such that  $x_j \neq x_{j+1}$  for any  $1 \leq j < k$ . We call  $k$  the *number of parts* and

$$d = 1 + \sum_{i=2}^k \begin{cases} 1 & \text{if } x_i \neq x_j \text{ for all } 1 \leq j < i, \\ 0 & \text{otherwise} \end{cases}$$

the *number of distinct part sizes*. The number  $a_c(n)$  of Carlitz compositions is [269]

$$a_c(n) \sim \frac{1}{\sigma F'(\sigma)} \left(\frac{1}{\sigma}\right)^n = (0.456387\dots)(1.750243\dots)^n$$

where  $\sigma = 0.571349\dots$  is the unique solution of the equation

$$F(x) = \sum_{j=1}^{\infty} (-1)^{j-1} \frac{x^j}{1-x^j} = 1, \quad 0 \leq x \leq 1.$$

The expected number of parts is asymptotically

$$\frac{G(\sigma)}{\sigma F'(\sigma)} n \sim (0.350571\dots)n \quad \text{where} \quad G(x) = \sum_{j=1}^{\infty} (-1)^{j-1} \frac{j x^j}{1-x^j}$$

(by contrast, an unrestricted composition has  $(n+1)/2$  parts on average). The expected size of the largest part is

$$\frac{-\ln(n)}{\ln(\sigma)} + \left( \frac{\ln(F'(\sigma)) + \ln(1-\sigma) - \gamma}{\ln(\sigma)} + \frac{1}{2} \right) + \varepsilon(n) = (1.786495\dots) \ln(n) + 0.64311\dots + \varepsilon(n)$$

where  $\gamma$  is Euler's constant and  $\varepsilon(n)$  is a small-amplitude zero-mean periodic function. The expected number of distinct part sizes is [270]

$$\frac{-\ln(n)}{\ln(\sigma)} + \left( \frac{\ln(F'(\sigma)) + \gamma}{\ln(\sigma)} + \frac{1}{2} \right) + \delta(n) = (1.786495\dots) \ln(n) - 2.932545\dots + \delta(n)$$

where  $\delta(n)$  is likewise negligible. (By contrast, an unrestricted composition has a largest part of size roughly  $\ln(n)/\ln(2) + 0.332746\dots$  and roughly  $\ln(n)/\ln(2) - 0.667253\dots$  distinct part sizes on average: see [271, 272, 273], as well as the bottom of page 340.) We wonder about the multiplicative analog of these results. See also [274].

**5.6. Otter's Tree Enumeration Constants.** Higher-order asymptotic series for  $T_n$ ,  $t_n$  and  $B_n$  are given in [89]. Also, the asymptotic analysis of series-parallel

posets [275] is similar to that of trees. See [276, 277] for more about  $k$ -gonal 2-trees, as well as a new formula for  $\alpha$  in terms of rational expressions involving  $e$ .

The generating function  $L(x)$  of leftist trees satisfies a simpler functional equation than previously thought:

$$L(x) = x + L(xL(x))$$

which involves an unusual nested construction. The radius of convergence  $\rho = 0.3637040915\dots = (2.7494879027\dots)^{-1}$  of  $L(x)$  satisfies

$$\rho L'(\rho L(\rho)) = 1$$

and the coefficient of  $\rho^{-n}n^{-1/2}$  in the asymptotic expression for  $L_n$  is

$$\sqrt{\frac{1}{2\pi\rho^2} \frac{\rho + L(\rho)}{L''(\rho L(\rho))}} = 0.2503634293\dots = (0.6883712204\dots)\rho.$$

The average height of  $n$ -leaf leftist trees is asymptotically  $(1.81349371\dots)\sqrt{\pi n}$  and the average depth of vertices belonging to such trees is asymptotically  $(0.90674685\dots)\sqrt{\pi n}$ . Nogueira [278] conjectured that the ratio of the two coefficients is exactly 2, but his only evidence is numerical (to over 1000 decimal digits).

For the following, we consider only unordered forests whose connected components are (strongly) ordered binary trees. Let  $F_n$  denote the number of such forests with  $2n - 1$  vertices; then the generating function

$$\Phi(x) = 1 + \sum_{n=1}^{\infty} F_n x^n = 1 + x + 2x^2 + 4x^3 + 10x^4 + 26x^5 + 77x^6 + \dots$$

satisfies

$$\Phi(x) = \exp\left(\sum_{k=1}^{\infty} \frac{1 - \sqrt{1 - 4x^k}}{2k}\right) = \prod_{m=1}^{\infty} (1 - x^m)^{-\frac{1}{m} \binom{2m-2}{m-1}}.$$

It can be shown that [265]

$$F_n \sim \frac{\Phi(1/4)}{\sqrt{\pi}} \frac{4^{n-1}}{n^{3/2}} = \frac{1.7160305349\dots}{4\sqrt{\pi}} \frac{4^n}{n^{3/2}}$$

as  $n \rightarrow \infty$ . The constant 1.716... also plays a role in the asymptotic analysis of the probability that a random forest has no two components of the same size.

**5.7. Lengyel's Constant.** Constants of the form  $\sum_{k=-\infty}^{\infty} 2^{-k^2}$  and  $\sum_{k=-\infty}^{\infty} 2^{-(k-1/2)^2}$  appear in [279, 280].

**5.9. Pólya's Random Walk Constants.** Properties of the gamma function lead to a further simplification [281]:

$$m_3 = \frac{1}{32\pi^3} (\sqrt{3} - 1) \left[ \Gamma\left(\frac{1}{24}\right) \Gamma\left(\frac{11}{24}\right) \right]^2$$

Consider a variation in which the drunkard performs a random walk starting from the origin with  $2^d$  equally probable steps, each of the form  $(\pm 1, \pm 1, \dots, \pm 1)$ . The number of walks that end at the origin after  $2n$  steps is

$$\tilde{U}_{d,0,2n} = \binom{2n}{n}^d$$

and the number of such walks for which  $2n$  is the time of *first return* to the origin is  $\tilde{V}_{d,0,2n}$ , where [282]

$$\begin{aligned} 2^{-n}\tilde{V}_{1,0,2n} &= \frac{1}{n2^{2n-1}} \binom{2n-2}{n-1} \sim \frac{1}{2\sqrt{\pi}n^{3/2}}, \\ 2^{-2n}\tilde{V}_{2,0,2n} &= \frac{\pi}{n(\ln(n))^2} - 2\pi \frac{\gamma + \pi B}{n(\ln(n))^3} + O\left(\frac{1}{n(\ln(n))^4}\right), \\ 2^{-3n}\tilde{V}_{3,0,2n} &= \frac{1}{\pi^{3/2}C^2n^{3/2}} + O\left(\frac{1}{n^2}\right) \end{aligned}$$

as  $n \rightarrow \infty$ , where

$$\begin{aligned} B &= 1 + \sum_{k=1}^{\infty} \left[ 2^{-4k} \binom{2k}{k}^2 - \frac{1}{\pi k} \right] = 0.8825424006\dots, \\ C &= \sum_{k=0}^{\infty} 2^{-6k} \binom{2k}{k}^3 = \frac{1}{4\pi^3} \Gamma\left(\frac{1}{4}\right)^4 = 1.3932039296\dots \end{aligned}$$

It turns out that the constant  $\sigma$ , given by an infinite series, has a more compact integral expression [283, 284]:

$$\sigma = \frac{1}{\pi} \int_0^{\infty} \frac{1}{x^2} \ln \left[ \frac{6}{x^2} \left( 1 - \frac{\sin(x)}{x} \right) \right] dx = -0.2979521902\dots = \frac{-0.5160683318\dots}{\sqrt{3}}$$

and surprisingly appears in both 3D statistical mechanics [285] and 1D probabilistic algorithms [286].

**5.10. Self-Avoiding Walk Constants.** Hueter [287, 288] rigorously proved that  $\nu_2 = 3/4$  and that  $7/12 \leq \nu_3 \leq 2/3$ ,  $1/2 \leq \nu_4 \leq 5/8$  (if the mean square end-to-end distance exponents  $\nu_3, \nu_4$  exist; otherwise the bounds apply for

$$\underline{\nu}_d = \liminf_{n \rightarrow \infty} \frac{\ln(r_n)}{2 \ln(n)}, \quad \bar{\nu}_d = \limsup_{n \rightarrow \infty} \frac{\ln(r_n)}{2 \ln(n)}$$

when  $d = 3, 4$ ). She confirmed that the same exponents apply for the mean square radius of gyration  $s_n$  for  $d = 2, 3, 4$ ; the results carry over to self-avoiding trails as

well [289]. Burkhardt & Guim [290] adjusted the estimate for  $\lim_{k \rightarrow \infty} p_{k,k}^{1/k^2}$  to 1.743...; this has now further been improved to 1.74455... [291].

**5.11. Feller's Coin Tossing Constants.** Additional references on oscillatory phenomena in probability theory include [292, 293, 294]; see also our earlier entry [5.5].

**5.12. Hard Square Entropy Constant.** McKay [295] observed the following asymptotic behavior:

$$F(n) \sim (1.06608266\dots)(1.0693545387\dots)^{2n}(1.5030480824\dots)^{n^2}$$

based on an analysis of the terms  $F(n)$  up to  $n = 19$ . He emphasized that the form of right hand side is conjectural, even though the data showed quite strong convergence to this form.

Let  $L(m, n)$  denote the number of legal positions on an  $m \times n$  Go board (a popular game). Then [296]

$$\begin{aligned} \lim_{n \rightarrow \infty} L(1, n)^{1/n} &= 1 + \frac{1}{3} \left( (27 + 3\sqrt{57})^{1/3} + (27 - 3\sqrt{57})^{1/3} \right) = 2.7692923542\dots, \\ \lim_{n \rightarrow \infty} L(n, n)^{1/n^2} &= 2.9757341920\dots \end{aligned}$$

and, subject to a plausible conjecture,

$$L(m, n) \sim (0.8506399258\dots)(0.96553505933\dots)^{m+n}(2.9757341920\dots)^{mn}$$

as  $\min\{m, n\} \rightarrow \infty$ .

**5.13. Binary Search Tree Constants.** The random permutation model for generating weakly binary trees (given an  $n$ -vector of distinct integers, construct  $T$  via insertions) does *not* provide equal weighting on the  $\binom{2n}{n}/(n+1)$  possible trees. For example, when  $n = 3$ , the permutations  $(2, 1, 3)$  and  $(2, 3, 1)$  both give rise to the same tree  $S$ , which hence has probability  $q(S) = 1/3$  whereas  $q(T) = 1/6$  for the other four trees. Fill [282, 297, 298] asked how the numbers  $q(T)$  themselves are distributed, for fixed  $n$ . If the trees are endowed with the uniform distribution, then

$$\begin{aligned} \frac{-\mathbb{E}[\ln(q(T))]}{n} &\rightarrow \sum_{k=1}^{\infty} \frac{\ln(k)}{(k+1)4^k} \binom{2k}{k} \\ &= -\gamma - \int_0^1 \frac{\ln(\ln(1/t))}{\sqrt{1-t}(1+\sqrt{1-t})^2} dt = 2.0254384677\dots \end{aligned}$$

as  $n \rightarrow \infty$ . If, instead, the trees follow the distribution  $q$ , then

$$\frac{-\mathbb{E}[\ln(q(T))]}{n} \rightarrow 2 \sum_{k=1}^{\infty} \frac{\ln(k)}{(k+1)(k+2)}$$



$$= -\gamma - 2 \int_0^1 \frac{((t-2)\ln(1-t) - 2t)\ln(\ln(1/t))}{t^3} dt = 1.2035649167\dots$$

The maximum value of  $-\ln(q(T))$  is  $\sim n \ln(n)$  and the minimum value is  $\sim cn$ , where

$$c = \ln(4) + \sum_{k=1}^{\infty} 2^{-k} \ln(1 - 2^{-k}) = 0.9457553021\dots$$

**5.14. Digital Search Tree Constants.** The constant  $Q$  is transcendental via a general theorem on values of modular forms due to Nesterenko [134, 135]. A correct formula for  $\theta$  is

$$\theta = \sum_{k=1}^{\infty} \frac{k2^{k(k-1)/2}}{1 \cdot 3 \cdot 7 \cdots (2^k - 1)} \sum_{j=1}^k \frac{1}{2^j - 1} = 7.7431319855\dots$$

(the exponent  $k(k-1)/2$  was mistakenly given as  $k+1$  in [299], but the numerical value is correct). The constants  $\alpha$ ,  $\beta$  and  $Q^{-1}$  appear in [300]. Also,  $\alpha$  appears in [301] and  $Q^{-1}$  appears in [280]. It turns out that  $\nu$  and  $\chi$  are linked via  $\nu - 1 = \chi$ ; we have [302, 303, 304]

$$\sum_{j=1}^{\infty} \frac{(-1)^{j-1}}{j(2^j - 1)} = \sum_{k=1}^{\infty} \ln(1 + 2^{-k}) = 0.8688766526\dots = \frac{7.2271128245\dots}{12 \ln(2)}.$$

Finally, a random variable  $X$  with density  $e^{-x}(e^{-x} - 1 + x)/(1 - e^{-x})^2$ ,  $x \geq 0$ , has mean  $E(X) = \pi^2/6$  and mean fractional part [304]

$$E(X - \lfloor X \rfloor) = \frac{11}{24} + \sum_{m=1}^{\infty} \frac{\pi^2}{\sinh(2\pi^2 m)^2} = \frac{11}{24} + (2.825535\dots) \times 10^{-16}.$$

The distribution of  $X$  is connected with the *random assignment problem* [305, 306].

**5.15. Optimal Stopping Constants.** When discussing the expected rank  $R_n$ , we assumed that no applicant would ever refuse a job offer! If each applicant only accepts an offer with known probability  $p$ , then [307]

$$\lim_{n \rightarrow \infty} R_n = \prod_{i=1}^{\infty} \left( 1 + \frac{2}{i} \frac{1 + pi}{2 - p + pi} \right)^{\frac{1}{1+pi}}$$

which is 6.2101994550... in the event that  $p = 1/2$ .

When discussing the full-information problem for Uniform  $[0, 1]$  variables, we assumed that the number of applicants is known. If instead this itself is a uniformly

distributed variable on  $\{1, 2, \dots, n\}$ , then for the “nothing but the best objective”, the asymptotic probability of success is [308, 309]

$$(1 - e^a) \operatorname{Ei}(-a) - (e^{-a} + a \operatorname{Ei}(-a))(\gamma + \ln(a) - \operatorname{Ei}(a)) = 0.4351708055\dots$$

where  $a = 2.1198244098\dots$  is the unique positive solution of the equation

$$e^a(1 - \gamma - \ln(a) + \operatorname{Ei}(-a)) - (\gamma + \ln(a) - \operatorname{Ei}(a)) = 1.$$

It is remarkable that these constants occur in other, seemingly unrelated versions of the secretary problem [310, 311, 312].

Suppose that you view successively terms of a sequence  $X_1, X_2, X_3, \dots$  of independent random variables with a common distribution function  $F$ . You know the function  $F$ , and as  $X_k$  is being viewed, you must either stop the process or continue. If you stop at time  $k$ , you receive a payoff  $(1/k) \sum_{j=1}^k X_j$ . Your objective is to maximize the expected payoff. An optimal strategy is to stop at the first  $k$  for which  $\sum_{j=1}^k X_j \geq \alpha_k$ , where  $\alpha_1, \alpha_2, \alpha_3, \dots$  are certain values depending on  $F$ . Shepp [313, 314] proved that  $\lim_{k \rightarrow \infty} \alpha_k / \sqrt{k}$  exists and is independent of  $F$  as long as  $F$  has zero mean and unit variance; further,

$$\lim_{k \rightarrow \infty} \frac{\alpha_k}{\sqrt{k}} = x = 0.8399236756\dots$$

is the unique zero of  $2x - \sqrt{2\pi} (1 - x^2) \exp(x^2/2) (1 + \operatorname{erf}(x/\sqrt{2}))$ . We wonder if Shepp’s constant can be employed to give a high-precision estimate of the Chow-Robbins constant  $2(0.7929535064\dots) - 1 = 0.5859070128\dots$  [315], the value of the expected payoff for  $F(-1) = F(1) = 1/2$ .

Also, consider a random binary string  $Y_1 Y_2 Y_3 \dots Y_n$  with  $P(Y_k = 1) = 1 - P(Y_k = 0)$  independent of  $k$  and  $Y_k$  independent of the other  $Y$ s. Let  $H$  denote the pattern consisting of the digits

$$\underbrace{1000\dots 0}_l \quad \text{or} \quad \underbrace{0111\dots 1}_l$$

and assume that its probability of occurrence for each  $k$  is

$$P(Y_{k+1} Y_{k+2} Y_{k+3} \dots Y_{k+l} = H) = \frac{1}{l} \left(1 - \frac{1}{l}\right)^{l-1} \sim \frac{1}{el} = \frac{0.3678794411\dots}{l}.$$

You observe sequentially the digits  $Y_1, Y_2, Y_3, \dots$  one at a time. You know the values  $n$  and  $l$ , and as  $Y_k$  is being observed, you must either stop the process or continue. Your objective is to stop at the final appearance of  $H$  up to  $Y_n$ . Bruss & Louchard [316] determined a strategy that maximizes the probability of meeting this goal. For  $n \geq \beta l$ , this success probability is

$$\frac{2}{135} e^{-\beta} (4 - 45\beta^2 + 45\beta^3) = 0.6192522709\dots$$

as  $l \rightarrow \infty$ , where  $\beta = 3.4049534663\dots$  is the largest zero of the cubic  $45\beta^3 - 180\beta^2 + 90\beta + 4$ . Further, the interval  $[0.367\dots, 0.619\dots]$  constitutes “typical” asymptotic bounds on success probabilities associated with a wide variety of optimal stopping problems in strings.

Suppose finally that you view a sequence  $Z_1, Z_2, \dots, Z_n$  of independent Uniform  $[0, 1]$  variables and that you wish to stop at a value of  $Z$  as large as possible. If you are a prophet (meaning that you have complete foresight), then you know  $Z_n^* = \max\{Z_1, \dots, Z_n\}$  beforehand and clearly  $E(Z_n^*) \sim 1 - 1/n$  as  $n \rightarrow \infty$ . If you are a 1-mortal (meaning that you have 1 opportunity to choose a  $Z$  via stopping rules) and if you proceed optimally, then the value  $Z_1^*$  obtained satisfies  $E(Z_1^*) \sim 1 - 2/n$ . If you are a 2-mortal (meaning that you have 2 opportunities to choose  $Z$ s and then take the maximum of these) and if you proceed optimally, then the value  $Z_2^*$  obtained satisfies  $E(Z_2^*) \sim 1 - c/n$ , where [317]

$$c = \frac{2\xi}{\xi + 2} = 1.1656232877\dots$$

and  $\xi = 2.7939976526\dots$  is the unique positive solution of the equation

$$\left(\frac{2}{\xi} + 1\right) \ln\left(\frac{\xi}{2} + 1\right) = \frac{3}{2}.$$

The performance improvement in having two choices over just one is impressive:  $c$  is much closer to 1 than 2!

**5.16. Extreme Value Constants.** The median of the Gumbel distribution is  $-\ln(\ln(2)) = 0.3665129205\dots$

**5.17. Pattern-Free Word Constants.** Recent references include [318, 319, 320].

**5.18. Percolation Cluster Density Constants.** An integral similar to that for  $\kappa_B(p_c)$  on the triangular lattice appears in [321]. Older references on 2D and 3D continuum percolation include [322, 323, 324, 325, 326]. See also [327, 328, 329].

Two infinite 0-1 sequences  $X, Y$  are called *compatible* if 0s can be deleted from  $X$  and/or from  $Y$  in such a way that the resulting 0-1 sequences  $X', Y'$  never have a 1 in the same position. For example, the sequences  $X = 000110\dots$  and  $Y = 110101\dots$  are not compatible. Assume that  $X$  and  $Y$  are randomly generated with each  $X_i, Y_j$  independent and  $P(X_i = 1) = P(Y_j = 1) = p$ . Intuition suggests that  $X$  and  $Y$  are compatible with positive probability if and only if  $p$  is suitably small. What is the supremum  $p^*$  of such  $p$ ? It is known [330, 331, 332, 333] that  $100^{-400} < p^* < 1/2$ ; simulation indicates [334] that  $0.3 < p^* < 0.305$ .

**5.19. Klarner’s Polyomino Constant.** A new estimate  $4.0625696\dots$  for  $\alpha$  is reported in [335] and a new rigorous lower bound of  $3.980137\dots$  in [336]. The number

$\bar{A}(n)$  of row-convex  $n$ -ominoes satisfies [337]

$$\bar{A}(n) = 5\bar{A}(n-1) - 7\bar{A}(n-2) + 4\bar{A}(n-3), \quad n \geq 5,$$

with  $\bar{A}(1) = 1$ ,  $\bar{A}(2) = 2$ ,  $\bar{A}(3) = 6$  and  $\bar{A}(4) = 19$ ; hence  $\bar{A}(n) \sim uv^n$  as  $n \rightarrow \infty$ , where  $v = 3.2055694304\dots$  is the unique real zero of  $x^3 - 5x^2 + 7x - 4$  and  $u = (41v^2 - 129v + 163)/944 = 0.1809155018\dots$

**5.20. Longest Subsequence Constants.** The Sankoff-Mainville conjecture that  $\lim_{k \rightarrow \infty} \gamma_k k^{1/2} = 2$  was proved by Kiwi, Loebel & Matousek [338]; the constant 2 arises from a connection with the longest increasing subsequence problem. A deeper connection with the Tracy-Widom distribution from random matrix theory has now been confirmed [339]:

$$\mathbb{E}(\lambda_{n,k}) \sim 2k^{-1/2}n + c_1 k^{-1/6} n^{1/3}, \quad \text{Var}(\lambda_{n,k}) \sim c_0 k^{-1/3} n^{2/3}$$

where  $k \rightarrow \infty$ ,  $n \rightarrow \infty$  in such a way that  $n/k^{1/2} \rightarrow 0$ .

Define  $\lambda_{n,k,r}$  to be the length of the longest common subsequence  $c$  of  $a$  and  $b$  subject to the constraint that, if  $a_i = b_j$  are paired when forming  $c$ , then  $|i - j| \leq r$ . Define as well  $\gamma_{k,r} = \lim_{n \rightarrow \infty} \mathbb{E}(\lambda_{n,k,r})/n$ . It is not surprising [340] that  $\lim_{r \rightarrow \infty} \gamma_{k,r} = \gamma_k$ . Also,  $\gamma_{2,1} = 7/10$ , but exact values for  $\gamma_{3,1}$ ,  $\gamma_{4,1}$ ,  $\gamma_{2,2}$  and  $\gamma_{2,3}$  remain open.

The Tracy-Widom distribution (specifically,  $F_{\text{GOE}}(x)$  as described in [341]) seems to play a role in other combinatorial problems [342, 343, 344], although the data is not conclusive. See also [345, 346, 347].

**5.21.  $k$ -Satisfiability Constants.** On the one hand, the lower bound for  $r_c(3)$  was improved to 3.42 in [348] and further improved to 3.52 in [349]. On the other hand, the upper bound 4.506 for  $r_c(3)$  in [350] has not been confirmed; the preceding two best upper bounds were 4.596 [351] and 4.571 [352].

**5.22. Lenz-Ising Constants.** The constant  $4G/\pi = 1.1662436161\dots$  appears in [353], as well as a new closed-form evaluation:

$$\begin{aligned} & \frac{\ln(2)}{2} + \frac{1}{16\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \ln[7 - 3\cos(\theta) - 3\cos(\varphi) - \cos(\theta)\cos(\varphi)] \, d\theta \, d\varphi \\ &= \frac{G}{\pi} + \frac{1}{2} \ln(\sqrt{2} - 1) + \frac{1}{\pi} \text{Ti}_2(3 + 2\sqrt{2}) = 0.7866842753\dots \end{aligned}$$

where  $\text{Ti}_2(x)$  is the inverse tangent integral (discussed on p. 57).

**5.23. Monomer-Dimer Constants.** Friedland & Peled [354] revisited Baxter's computation of  $A$  and confirmed that  $\ln(A) = 0.66279897\dots$ . They examined the three-dimensional analog,  $A'$ , of  $A$  and found that  $0.7652 < \ln(A') < 0.7863$ . See also [355, 356].

**5.25. Tutte-Beraha Constants.** For any positive integer  $r$ , there is a best constant  $C(r)$  such that, for each graph of maximum degree  $\leq r$ , the complex zeros of its chromatic polynomial lie in the disk  $|z| \leq C(r)$ . Further,  $K = \lim_{r \rightarrow \infty} C(r)/r$  exists and  $K = 7.963906\dots$  is the smallest number for which

$$\inf_{\alpha > 0} \frac{1}{\alpha} \sum_{n=2}^{\infty} e^{\alpha n} K^{-(n-1)} \frac{n^{n-1}}{n!} \leq 1.$$

Sokal [357] proved all of the above, answering questions raised in [358, 359]. See also [360].

**6.1. Gauss' Lemniscate Constant.** Consider the following game [361]. Players  $A$  and  $B$  simultaneously choose numbers  $x$  and  $y$  in the unit interval;  $B$  then pays  $A$  the amount  $|x - y|^{1/2}$ . The value of the game (that is, the expected payoff, assuming both players adopt optimal strategies) is  $M/2 = 0.59907\dots$ . Also, let  $\xi_1, \xi_2, \dots, \xi_n, \eta_1, \eta_2, \dots, \eta_n$  be distinct points in the plane and construct, with these points as centers, squares of side  $s$  and of arbitrary orientation that do not overlap. Then

$$s \leq \frac{L}{\sqrt{2}} \left( \frac{\prod_{i=1}^n \prod_{j=1}^n |\xi_i - \eta_j|}{\prod_{i < j} |\xi_i - \xi_j| \cdot \prod_{i < j} |\eta_i - \eta_j|} \right)^{1/n}$$

and the constant  $L/\sqrt{2} = 1.85407\dots$  is best possible [362].

**6.2. Euler-Gompertz Constant.** The two quantities

$$I_0(2) = \sum_{k=0}^{\infty} \frac{1}{(k!)^2} = 2.2795853023\dots, \quad J_0(2) = \sum_{k=0}^{\infty} \frac{(-1)^k}{(k!)^2} = 0.2238907791\dots$$

are similar, but only the first is associated with continued fractions. Here is an interesting occurrence of the second: letting [363]

$$a_0 = a_1 = 1, \quad a_n = n a_{n-1} - a_{n-2} \quad \text{for } n \geq 2,$$

we have  $\lim_{n \rightarrow \infty} a_n/n! = J_0(2)$ .

**6.3. Kepler-Bouwkamp Constant.** Additional references include [364, 365, 366] and another representation is [367]

$$\rho = \frac{3^{10} \sqrt{3}}{2^7 5^2 7 11 \pi} \exp \left[ - \sum_{k=1}^{\infty} \frac{(\zeta(2k) - 1 - 2^{-2k} - 3^{-2k}) 2^{2k} (\lambda(2k) - 1 - 3^{-2k})}{k} \right];$$

the series converges at the same rate as a geometric series with ratio  $1/100$ . A relevant inequality is [368]

$$\int_0^{\infty} \cos(2x) \prod_{j=1}^{\infty} \cos\left(\frac{x}{j}\right) dx < \frac{\pi}{8}$$

and the difference is less than  $10^{-42}$ ! A prime analog of  $\rho$  is [369]

$$\prod_{p \geq 3} \cos\left(\frac{\pi}{p}\right) = 0.3128329295\dots = (3.1965944300\dots)^{-1}$$

and variations abound.

**6.5. Plouffe's Constant.** This constant is included in a fascinating mix of ideas by Smith [370], who claims that “angle-doubling” one bit at a time was known centuries ago to Archimedes and was implemented decades ago in binary cordic algorithms (also mentioned in section 5.14). Another constant of interest is  $\arctan(\sqrt{2}) = 0.9553166181\dots$ , which is the base angle of a certain isosceles spherical triangle (in fact, the unique non-Euclidean triangle with rational sides and a single right angle).

Chowdhury [371] generalized his earlier work on bitwise XOR sums and the logistic map: A sample new result is

$$\sum_{n=0}^{\infty} \frac{\rho(b_n b_{n-1})}{2^{n+1}} = \frac{1}{4\pi} \oplus \frac{1}{\pi}$$

where  $b_n = \cos(2^n)$ . The right-hand side is computed merely by shifting the binary expansion of  $1/\pi$  two places (to obtain  $1/(4\pi)$ ) and adding modulo two without carries (to find the sum).

**6.6. Lehmer's Constant.** Rivoal [372] has studied the link between the rational approximations of a positive real number  $x$  coming from the continued cotangent representation of  $x$ , and the usual convergents that proceed from the regular continued fraction expansion of  $x$ .

**6.8. Prouhet-Thue-Morse Constant.** Simple analogs of the Woods-Robbins and Flajolet-Martin formulas are [47]

$$\prod_{m=1}^{\infty} \left(\frac{2m}{2m-1}\right)^{(-1)^m} = \frac{\sqrt{2}\pi^{3/2}}{\Gamma(1/4)^2}, \quad \prod_{m=1}^{\infty} \left(\frac{2m}{2m+1}\right)^{(-1)^m} = \frac{\Gamma(1/4)^2}{2^{5/2}\sqrt{\pi}};$$

we wonder about the outcome of exponent sequences other than  $(-1)^m$  or  $(-1)^{t_m}$ .

**6.9. Minkowski-Bower Constant.** See [373, 374] for a generalization of the Minkowski question mark function.

**6.10. Quadratic Recurrence Constants.** The sequence  $k_{n+1} = (1/n)k_n^2$ , where  $n \geq 0$ , is convergent if and only if

$$|k_0| < \exp\left(\sum_{j=2}^{\infty} \frac{\ln(j)}{2^j}\right) = 1.6616879496\dots$$

Moreover, the sequence either converges to zero or diverges to infinity [375, 376]. A systematic study of threshold constants like this, over a broad class of quadratic recurrences, has never been attempted. The constant 1.2640847353... and Sylvester's sequence appear in an algebraic-geometric setting [377]. Also, results on Somos' sequences are found in [378, 379] and on the products

$$1^{1/2}2^{1/4}3^{1/8}\dots = 1.6616879496\dots, \quad 1^{1/3}2^{1/9}3^{1/27}\dots = 1.1563626843\dots$$

in [47, 68].

**6.11. Iterated Exponential Constants.** Consider the recursion

$$a_1 = 1, \quad a_n = a_{n-1} \exp\left(\frac{1}{ea_{n-1}}\right)$$

for  $n \geq 2$ . It is known that [380]

$$a_n = \frac{n}{e} + \frac{\ln(n)}{2e} + \frac{C}{e} + o(1), \quad (n!)^{1/n} = \frac{n}{e} + \frac{\ln(n)}{2e} + \frac{\ln(\sqrt{2\pi})}{e} + o(1)$$

as  $n \rightarrow \infty$ , where

$$C = e - 1 + \frac{\gamma}{2} + \frac{1}{2} \sum_{k=1}^{\infty} \frac{k - ea_k}{eka_k} + \sum_{k=1}^{\infty} \left( ea_{k+1} - ea_k - 1 - \frac{1}{2ea_k} \right) = 1.2905502\dots$$

Further,  $a_n - (n!)^{1/n}$  is strictly increasing and

$$a_n - (n!)^{1/n} \leq (C - \ln(\sqrt{2\pi}))/e = 0.136708\dots$$

for all  $n$ . The constant is best possible. Putting  $b_n = 1/(ea_n)$  yields the recursion  $b_n = b_{n-1} \exp(-b_{n-1})$ , for which an analogous asymptotic expansion can be written.

The unique real zero  $z_n$  of  $\sum_{k=0}^n z^k/k!$ , where  $n$  is odd, satisfies  $\lim_{n \rightarrow \infty} z_n/n = W(e^{-1}) = 0.2784645427\dots$  [381, 382]. Also,  $3^{-1}e^{-1/3} = 0.2388437701\dots$  arises in [383] as a consequence of the formula  $-W(-3^{-1}e^{-1/3}) = 1/3$ . Note that  $-W(-x)$  is the exponential generating function for rooted labeled trees and hence is often called the *tree function*.

**6.12. Conway's Constant.** A "biochemistry" based on Conway's "chemistry" appears in [384].

**7.1. Bloch-Landau Constants.** In the definitions of the sets  $F$  and  $G$ , the functions  $f$  need only be analytic on the open unit disk  $D$  (in addition to satisfying  $f(0) = 0$ ,  $f'(0) = 1$ ). On the one hand, the weakened hypothesis doesn't affect the values of  $B$ ,  $L$  or  $A$ ; on the other hand, the weakening is essential for the existence of  $f \in G$  such that  $m(f) = M$ .

The bounds  $0.62\pi < A < 0.7728\pi$  were improved by several authors, although they studied the quantity  $\tilde{A} = \pi - A$  instead (the omitted area constant). Barnard & Lewis [385] demonstrated that  $\tilde{A} \leq 0.31\pi$ . Barnard & Pearce [386] established that  $\tilde{A} \geq 0.240005\pi$ , but Banjai & Trefethen [387] subsequently computed that  $\tilde{A} = (0.2385813248\dots)\pi$ . It is believed that the earlier estimate was slightly in error. See [388, 389, 390] for related problems.

The spherical analog of Bloch's constant  $B$ , corresponding to meromorphic functions  $f$  mapping  $D$  to the Riemann sphere, was recently determined by Bonk & Eremenko [391]. This constant turns out to be  $\arccos(1/3) = 1.2309594173\dots$ . A proof as such gives us hope that someday the planar Bloch-Landau constants will also be exactly known.

More relevant material is found in [226, 392].

**7.2. Masser-Gramain Constant.** Suppose  $f(z)$  is an entire function such that  $f^{(k)}(n)$  is an integer for each nonnegative integer  $n$ , for each integer  $0 \leq k \leq s - 1$ . (We have discussed only the case  $s = 1$ .) The best constant  $\theta_s > 0$  for which

$$\limsup_{r \rightarrow \infty} \frac{\ln(M_r)}{r} < \theta_s \quad \text{implies } f \text{ is a polynomial}$$

was proved by Bundschuh & Zudilin [393], building on Gel'fond [394] and Selberg [395], to satisfy

$$s \cdot \frac{\pi}{3} \geq \theta_s > \begin{cases} 0.994077\dots & \text{if } s = 2, \\ 1.339905\dots & \text{if } s = 3, \\ 1.674474\dots & \text{if } s = 4. \end{cases}$$

(Actually they proved much more.) Can a Gaussian integer-valued analog of these integer-valued results be found?

**7.5. Hayman Constants.** A new upper bound [396] for the Hayman-Korenblum constant  $c(2)$  is 0.69472. An update on the Hayman-Wu constant appears in [397].

**7.6. Littlewood-Clunie-Pommerenke Constants.** The lower limit of summation in the definition of  $S_2$  should be  $n = 0$  rather than  $n = 1$ , that is, the coefficient  $b_0$  need not be zero. We have sharp bounds  $|b_1| \leq 1$ ,  $|b_2| \leq 2/3$ ,  $|b_3| \leq 1/2 + e^{-6}$  [398]. The bounds on  $\gamma_k$  due to Clunie & Pommerenke should be 0.509 and 0.83 [399]; Carleson & Jones' improvement was nonrigorous. While  $0.83 = 1 - 0.17$  remains the best established upper bound, the lower bound has been increased to  $0.54 = 1 - 0.46$  [400, 401, 402]. Numerical evidence for both the Carleson-Jones conjecture and Brennan's conjecture was found by Kraetzer [403]. Theoretical evidence supporting the



latter appears in [404], but a complete proof remains undiscovered. It seems that  $\alpha = 1 - \gamma$  is now a theorem [405, 406] whose confirmation is based on the recent work of several researchers [407, 408, 409].

**8.1. Geometric Probability Constants.** Just as the ratio of a semicircle to its diameter is always  $\pi/2$ , the ratio of the latus rectum arc of any parabola to its latus rectum is [410, 411]

$$\frac{1}{2} \left( \sqrt{2} + \ln(1 + \sqrt{2}) \right) = 1.1477935746\dots = \frac{1}{2} (2.2955871493\dots)$$

Is it mere coincidence that this constant is so closely related to the quantity  $\delta(2)$ ? The expected distance between two random points on different sides of the unit square is [368]

$$\frac{2 + \sqrt{2} + 5 \ln(1 + \sqrt{2})}{9} = 0.8690090552\dots$$

and the expected distance between two random points on different faces of the unit cube is

$$\frac{4 + 17\sqrt{2} - 6\sqrt{3} - 7\pi + 21 \ln(1 + \sqrt{2}) + 21 \ln(7 + 4\sqrt{3})}{75} = 0.9263900551\dots$$

Also, the convex hull of random point sets in the unit disk (rather than the unit square) is mentioned in [412].

**8.4. Moser's Worm Constant.** Relevant progress is described in [413]. We mention, in Figure 8.3, that the quantity  $x = \sec(\varphi) = 1.0435901095\dots$  is algebraic of degree six [415]:

$$3x^6 + 36x^4 + 16x^2 - 64 = 0$$

and wonder if this is linked to Figure 8.7 and the Reuleaux triangle of width 1.5449417003... (also algebraic of degree six [416]). The latter is the planar set of maximal constant width that avoids all vertices of the integer square lattice.

**8.5. Traveling Salesman Constants.** The cavity method is applied in [417] to matchings on sparse random graphs.

**8.7. Hermite's Constants.** A lattice  $\Lambda$  in  $\mathbb{R}^n$  consists of all integer linear combinations of a set of basis vectors  $\{e_j\}_{j=1}^n$  for  $\mathbb{R}^n$ . If the fundamental parallelepiped determined by  $\{e_j\}_{j=1}^n$  has Lebesgue measure 1, then  $\Lambda$  is said to be of unit volume. The constants  $\gamma_n$  can be defined via an optimization problem

$$\gamma_n = \max_{\substack{\text{unit volume} \\ \text{lattices } \Lambda}} \min_{\substack{x \in \Lambda, \\ x \neq 0}} \|x\|^2$$

and are listed in Table 8.10. The precise value of the next constant  $2 \leq \gamma_9 < 2.1327$  remains open [418, 419, 420], although Cohn & Kumar [421, 422] have recently proved

that  $\gamma_{24} = 4$ . A classical theorem [423, 424, 425] provides that  $\gamma_n^n$  is rational for all  $n$ . It is not known if the sequence  $\gamma_1, \gamma_2, \gamma_3, \dots$  is strictly increasing, or if the ratio  $\gamma_n/n$  tends to a limit as  $n \rightarrow \infty$ . See also [426, 427].

Table 8.10 *Hermite's Constants*  $\gamma_n$ 

$n$	Exact	Decimal	$n$	Exact	Decimal
1	1	1	5	$8^{1/5}$	1.5157165665...
2	$(4/3)^{1/2}$	1.1547005383...	6	$(64/3)^{1/6}$	1.6653663553...
3	$2^{1/3}$	1.2599210498...	7	$64^{1/7}$	1.8114473285...
4	$4^{1/4}$	1.4142135623...	8	2	2

An arbitrary packing of the plane with disks is called compact if every disk  $D$  is tangent to a sequence of disks  $D_1, D_2, \dots, D_n$  such that  $D_i$  is tangent to  $D_{i+1}$  for  $i = 1, 2, \dots, n$  with  $D_{n+1} = D$ . If we pack the plane using disks of radius 1, then the only possible compact packing is the hexagonal lattice packing with density  $\pi/\sqrt{12}$ . If we pack the plane using disks of radius 1 and  $r < 1$  (disks of both sizes must be used), then there are precisely nine values of  $r$  for which a compact packing exists. See Table 8.11. For seven of these nine values, it is known that the densest packing is a compact packing; the same is expected to be true for the remaining two values [428, 429, 430].

Table 8.11 *All Nine Values of  $r < 1$  Which Allow Compact Packings*

Exact (expression or minimal polynomial)	Decimal
$5 - 2\sqrt{6}$	0.1010205144...
$(2\sqrt{3} - 3)/3$	0.1547005383...
$(\sqrt{17} - 3)/4$	0.2807764064...
$x^4 - 28x^3 - 10x^2 + 4x + 1$	0.3491981862...
$9x^4 - 12x^3 - 26x^2 - 12x + 9$	0.3861061048...
$\sqrt{2} - 1$	0.4142135623...
$8x^3 + 3x^2 - 2x - 1$	0.5332964166...
$x^8 - 8x^7 - 44x^6 - 232x^5 - 482x^4 - 24x^3 + 388x^2 - 120x + 9$	0.5451510421...
$x^4 - 10x^2 - 8x + 9$	0.6375559772...

There is space to only mention the circle-packing rigidity constants  $s_n$  [414], their limiting behavior:

$$\lim_{n \rightarrow \infty} n s_n = \frac{2^{4/3} \Gamma(1/3)^2}{3 \Gamma(2/3)} = 4.4516506980\dots$$

and their connection with conformal mappings.

**8.10. Reuleaux Triangle Constants.** In our earlier entry [8.4], we ask about the connection between two relevant algebraic quantities [415, 416], both zeroes of sextic polynomials.

**8.15. Graham’s Hexagon Constant.** Bieri [431] partially anticipated Graham’s result. A nice presentation of Reinhardt’s isodiametric theorem is found in [432].

**8.18. Rectilinear Crossing Constant.** It is now known [433, 434, 435, 436, 437, 438, 439] that

$$\bar{\nu}(K_{13}) = 229, \quad \bar{\nu}(K_{14}) = 324, \quad \bar{\nu}(K_{15}) = 447, \quad \bar{\nu}(K_{16}) = 603, \quad \bar{\nu}(K_{17}) = 798$$

and  $0.37968 < \rho < 0.38058$ .

**8.19. Circumradius-Inradius Constants.** The phrase “ $Z$ -admissible” in the caption of Figure 8.22 should be replaced by “ $Z$ -allowable”.

**Table of Constants.** The formula corresponding to  $0.8427659133\dots$  is  $(12 \ln(2))/\pi^2$  and the formula corresponding to  $0.8472130848\dots$  is  $M/\sqrt{2}$ .

#### REFERENCES

- [1] T. M. Apostol, Irrationality of the square root of 2 – a geometric proof, *Amer. Math. Monthly* 107 (2000) 841–842.
- [2] R. L. Graham and H. O. Pollak, Note on a nonlinear recurrence related to  $\sqrt{2}$ , *Math. Mag.* 43 (1970) 143–145; MR0265269 (42 #180).
- [3] D. Shanks, Incredible identities, *Fibonacci Quart.* 12 (1974) 271, 280; W. G. Spohn, letter to the editor, 14 (1976) 12; MR0349623 (50 #2116) and MR0384744 (52 #5617).
- [4] A. Borodin, R. Fagin, J. E. Hopcroft and M. Tompa, Decreasing the nesting depth of expressions involving square roots, *J. Symbolic Comput.* 1 (1985) 169–188; MR0818877 (87a:68087).
- [5] R. Zippel, Simplification of expressions involving radicals, *J. Symbolic Comput.* 1 (1985) 189–210; MR0818878 (87b:68058).
- [6] S. Landau, A note on “Zippel denesting”, *J. Symbolic Comput.* 13 (1992) 41–45; MR1153633 (93b:11168).
- [7] C. Ciamberlini and A. Marengoni, Su una interessante curiosita numerica, *Periodico di Matematiche* 17 (1937) 25–30.
- [8] M. Lotan, A problem in difference sets, *Amer. Math. Monthly* 56 (1949) 535–541; MR0032553 (11,306h).

- [9] W. A. Webb, The length of the four-number game, *Fibonacci Quart.* 20 (1982) 33–35; MR0660757 (84e:10017).
- [10] J. H. Selleck, Powers of  $T$  and Soddy circles, *Fibonacci Quart.* 21 (1983) 250–252; MR0723783 (85f:11014).
- [11] D. G. Rogers, Malfatti’s problem for apprentice masons (and geometers), preprint (2004).
- [12] F. A. Potra, On an iterative algorithm of order 1.839... for solving nonlinear operator equations, *Numer. Funct. Anal. Optim.* 7 (1984/85) 75–106; MR0772168 (86j:47088).
- [13] A. Herschfeld, On infinite radicals, *Amer. Math. Monthly* 42 (1935) 419–429.
- [14] S. R. Finch, Optimal escape paths, unpublished note (2005).
- [15] E. Makover and J. McGowan, An elementary proof that random Fibonacci sequences grow exponentially, math.NT/0510159.
- [16] A. R. Kitson, The regularized product of the Fibonacci numbers, math.NT/0608187.
- [17] N. Pippenger, An infinite product for  $e$ , *Amer. Math. Monthly* 87 (1980) 391.
- [18] C. S. Davis, On some simple continued fractions connected with  $e$ , *J. London Math. Soc.* 20 (1945) 194–198; MR0017394 (8,148b).
- [19] R. F. C. Walters, Alternative derivation of some regular continued fractions, *J. Austral. Math. Soc.* 8 (1968) 205–212; MR0226245 (37 #1835).
- [20] K. R. Matthews and R. F. C. Walters, Some properties of the continued fraction expansion of  $(m/n)e^{1/q}$ , *Proc. Cambridge Philos. Soc.* 67 (1970) 67–74; MR0252889 (40 #6104).
- [21] A. J. van der Poorten, Continued fraction expansions of values of the exponential function and related fun with continued fractions, *Nieuw Arch. Wisk.* 14 (1996) 221–230; MR1402843 (97f:11011).
- [22] G. Martin, Denser Egyptian fractions, *Acta Arith.* 95 (2000) 231–260; MR1793163 (2001m:11040).
- [23] E. S. Croot, On unit fractions with denominators in short intervals, *Acta Arith.* 99 (2001) 99–114; MR1847616 (2002e:11045).

- [24] A. Levin, A new class of infinite products generalizing Viète's product formula for  $\pi$ , *Ramanujan J.* 10 (2005) 305–324; available online at <http://www.math.brown.edu/~adlevin/>; MR2193382 (2006h:11142).
- [25] A. Levin, A geometric interpretation of an infinite product for the lemniscate constant, *Amer. Math. Monthly* 113 (2006) 510–520; available online at <http://www.math.brown.edu/~adlevin/>.
- [26] C. L. Adler and J. Tanton,  $\pi$  is the minimum value for pi, *College Math. J.* 31 (2000) 102–106; MR1766159 (2001c:11141).
- [27] L. D. Servi, Nested square roots of 2, *Amer. Math. Monthly* 110 (2003) 326–330.
- [28] M. A. Nyblom, More nested square roots of 2, *Amer. Math. Monthly* 112 (2005) 822–825.
- [29] P. Kiss and F. Mátyás, An asymptotic formula for  $\pi$ , *J. Number Theory* 31 (1989) 255–259; MR0993902 (90e:11036).
- [30] B. Tropak, Some asymptotic properties of Lucas numbers, *Proc. Regional Mathematical Conf.*, Kalsk, 1988, ed. A. Grytczuk, Pedagog. Univ. Zielona Góra, 1990, pp. 49–55; MR1114366 (92e:11013).
- [31] S. Akiyama, Lehmer numbers and an asymptotic formula for  $\pi$ , *J. Number Theory* 36 (1990) 328–331; MR1077711 (91m:11005).
- [32] S. Akiyama, A new type of inclusion exclusion principle for sequences and asymptotic formulas for  $\zeta(k)$ , *J. Number Theory* 45 (1993) 200–214; MR1242715 (94k:11027).
- [33] J.-P. Bézivin, Plus petit commun multiple des termes consécutifs d'une suite récurrente linéaire, *Collect. Math.* 40 (1989) 1–11 (1990); MR1078087 (92f:11022).
- [34] J. Gibbons, Unbounded spigot algorithms for the digits of pi, *Amer. Math. Monthly* 113 (2006) 318–328; available online at <http://web.comlab.ox.ac.uk/oucl/work/jeremy.gibbons/>.
- [35] D. H. Bailey, A compendium of BBP-type formulas for mathematical constants, available online at <http://crd.lbl.gov/~dhbailey/dhbpapers/>.
- [36] H.-C. Chan, More formulas for  $\pi$ , *Amer. Math. Monthly* 113 (2006) 452–455; MR2225478 (2006m:11184).

- [37] P. G. Lejeune Dirichlet, Über die Bestimmung der mittleren Werthe in der Zahlentheorie, *Abhandlungen der Königlich-Preussischen Akademie der Wissenschaften in Berlin* (1849) 69–83; *Werke*, v. 2, ed. L. Kronecker and L. Fuchs, Reimer, 1897, pp. 49–66.
- [38] L. E. Dickson, *History of the Theory of Numbers*, v. 1, *Divisibility and Primality*, Chelsea, 1971; p. 327; MR0245499 (39 #6807a).
- [39] S. Krämer, Die Eulersche Konstante  $\gamma$  und verwandte Zahlen, Diplomarbeit, Georg-August Universität Göttingen, 2005.
- [40] N. Nielsen, Een Raekke for Euler’s Konstant, *Nyt Tidsskrift for Matematik* 8B (1897) 10–12; JFM 28.0235.02.
- [41] E. Jacobsthal, Über die Eulersche Konstante, *Math.-Naturwiss. Blätter* 9 (1906) 153–154; *Norske Vid. Selsk. Skr. (Trondheim)* (1967) 1–24; MR0231776 (38 #104).
- [42] S. Selberg, Über die Reihe für die Eulersche Konstante, die von E. Jacobsthal und V. Brun angegeben ist, *Norske Vid. Selsk. Forh.* 12 (1939) 89–92; MR0001315 (1,216e).
- [43] M. Koecher, Einige Bemerkungen zur Eulerschen Konstanten, *Bayer. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. 1989* (1990) 9–16; MR1086006 (91k:11118).
- [44] J. W. L. Glaisher, On Dr. Vacca’s series for  $\gamma$ , *Quart. J. Pure Appl. Math.* 41 (1910) 365–368.
- [45] J. Sondow, Double integrals for Euler’s constant and  $\ln(4/\pi)$  and an analog of Hadjicostas’s formula, *Amer. Math. Monthly* 112 (2005) 61–65; math.CA/0211148; MR2110113 (2005i:11181).
- [46] J. Sondow, New Vacca-type rational series for Euler’s constant and its “alternating” analog  $\ln(4/\pi)$ , math.NT/0508042.
- [47] J. Sondow and P. Hadjicostas, The generalized-Euler-constant function  $\gamma(z)$  and a generalization of Somos’s quadratic recurrence constant, math.CA/0610499.
- [48] J. Sondow, Criteria for irrationality of Euler’s constant, *Proc. Amer. Math. Soc.* 131 (2003) 3335–3344; math.NT/0209070; MR1990621 (2004b:11102).
- [49] J. Sondow, A hypergeometric approach, via linear forms involving logarithms, to irrationality criteria for Euler’s constant, math.NT/0211075.

- [50] J. Sondow and W. Zudilin, Euler's constant,  $q$ -logarithms, and formulas of Ramanujan and Gosper, math.NT/0304021.
- [51] T. Hessami Pilehrood and Kh. Hessami Pilehrood, Criteria for irrationality of generalized Euler's constant, *J. Number Theory* 108 (2004) 169–185; MR2078662 (2005e:11095).
- [52] M. Prévost, A family of criteria for irrationality of Euler's constant, math.NT/0507231.
- [53] K. Mahler, Applications of a theorem by A. B. Shidlovski, *Proc. Royal Soc. Ser. A* 305 (1968) 149–173; available online at <http://www.cecm.sfu.ca/Mahler/>; MR0225729 (37 #1322).
- [54] A. J. van der Poorten, Obituary: Kurt Mahler, 1903–1988, *J. Austral. Math. Soc. Ser. A* 51 (1991) 343–380; available online at <http://www.cecm.sfu.ca/Mahler/>; MR1125440 (93a:01055).
- [55] H. G. Diamond, A number theoretic series of I. Kasara, *Pacific J. Math.* 111 (1984) 283–285; MR0734855 (85c:11078).
- [56] W.-B. Zhang, On a number theoretic series, *J. Number Theory* 30 (1988) 109–119; MR0961910 (90b:11092).
- [57] B. S. Rimskiĭ-Korsakov, A version of the construction of a theory of generalized gamma-functions based on the Laplace transform (in Russian), *Moskov. Oblast. Pedagog. Inst. Uč. Zap.* 57 (1957) 121–141; Engl. transl. in *Amer. Math. Soc. Transl.* (2) 17 (1961) 201–217; MR0098851 (20 #5303) and MR0124535 (23 #A1847).
- [58] A. A. Markoff, Mémoire sur la transformation des séries peu convergentes en séries très convergentes, *Mémoires de L'Académie Impériale des Sciences de St.-Pétersbourg*, VII<sup>e</sup> série, t. 37, n. 9; available online at <http://www.math.mun.ca/~sergey/Research/History/Markov/markov1890.html>.
- [59] M. Kondratieva and S. Sadov, Markov's transformation of series and the WZ method, *Adv. Appl. Math.* 34 (2005) 393–407; math.CA/0405592; MR2110559 (2005k:65006).
- [60] M. Mohammed and D. Zeilberger, The Markov-WZ method (2004), available online at <http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/marwz.html>.
- [61] B. C. Berndt, *Ramanujan's Notebooks. Part II*, Springer-Verlag, 1989, p. 293; MR0970033 (90b:01039).

- [62] L. Vepstas, On Plouffe's Ramanujan identities, math.NT/0609775.
- [63] S. Plouffe, Identities inspired by Ramanujan notebooks (part 2), available online at <http://www.lacim.uqam.ca/~plouffe/>.
- [64] T. Rivoal and W. Zudilin, Diophantine properties of numbers related to Catalan's constant, *Math. Annalen* 326 (2003) 705–721; MR2003449 (2004k:11119).
- [65] F. Beukers, A note on the irrationality of  $\zeta(2)$  and  $\zeta(3)$ , *Bull. London Math. Soc.* 11 (1979) 268–272; MR0554391 (81j:10045).
- [66] V. S. Adamchik, Integral and series representations for Catalan's constant, entry 18, available online at <http://www-2.cs.cmu.edu/~adamchik/articles/catalan.htm>.
- [67] W. Zudilin, An Apéry-like difference equation for Catalan's constant, *Elec. J. Combin.* 10 (2003) R14; math.NT/0201024; MR1975764 (2004d:11125).
- [68] J. Guillera and J. Sondow, Double integrals and infinite products for some classical constants via analytic continuations of Lerch's transcendent, math.NT/0506319.
- [69] S. R. Finch, Class number theory, unpublished note (2005).
- [70] G. H. Choe, Generalized continued fractions, *Appl. Math. Comput.* 109 (2000) 287–299; MR1738196 (2000k:28028).
- [71] G. H. Choe and C. Kim, The Khintchine constants for generalized continued fractions, *Appl. Math. Comput.* 144 (2003) 397–411; MR1994080 (2004e:37013).
- [72] S. Fischler and T. Rivoal, Un exposant de densité en approximation rationnelle, preprint (2006), available online at <http://www-fourier.ujf-grenoble.fr/~rivoal/>.
- [73] P. Grassberger, On the Hausdorff dimension of fractal attractors, *J. Stat. Phys.* 26 (1981) 173–179; MR0643707 (83i:58063).
- [74] G. Levin and F. Przytycki, On Hausdorff dimension of some Cantor attractors, math.DS/0309191.
- [75] T. Ord and T. D. Kieu, On the existence of a new family of Diophantine equations for  $\Omega$ , *Fund. Inform.* 56 (2003) 273–284; math.NT/0301274; MR2024745 (2004m:03166).
- [76] Y. V. Matiyasevich, private communication (2005).



- [77] J. P. Jones, D. Sato, H. Wada and D. Wiens, Diophantine representation of the set of prime numbers, *Amer. Math. Monthly* 83 (1976) 449–464; MR0414514 (54 #2615).
- [78] C. S. Calude and M. A. Stay, From Heisenberg to Gödel via Chaitin, CDMTCS report 95, <http://www.cs.auckland.ac.nz/CDMTCS/researchreports/235cris.pdf>.
- [79] B. Green, Generalising the Hardy-Littlewood method for primes, math.NT/0601211.
- [80] J.-L. Nicolas, Sur la distribution des nombres entiers ayant une quantité fixée de facteurs premiers, *Acta Arith.* 44 (1984) 191–200; MR0774099 (86c:11067).
- [81] H.-K. Hwang, Sur la répartition des valeurs des fonctions arithmétiques. Le nombre de facteurs premiers d'un entier, *J. Number Theory* 69 (1998) 135–152; MR1618482 (99d:11100).
- [82] N. P. Romanoff, Über einige Sätze der additiven Zahlentheorie, *Math. Annalen* 57 (1934) 668–678.
- [83] P. Erdős, On integers of the form  $2^k + p$  and some related problems, *Summa Brasil. Math.* 2 (1950) 113–123; MR0044558 (13,437i).
- [84] Y.-G. Chen and X.-G. Sun, On Romanoff's constant, *J. Number Theory* 106 (2004) 275–284; MR2059075 (2005g:11195).
- [85] G. H. Hardy and S. Ramanujan, Asymptotic formulae for the distribution of integers of various types, *Proc. London Math. Soc.* 16 (1917) 112–132; also in *Collected Papers of G. H. Hardy*, v. 1, Oxford Univ. Press, 1966, pp. 277–293.
- [86] L. B. Richmond, Asymptotic results for partitions. I. The distribution of certain integers, *J. Number Theory* 8 (1976) 372–389; MR0429806 (55 #2816).
- [87] J.-M. De Koninck and G. Tenenbaum, Sur la loi de répartition du  $k$ -ième facteur premier d'un entier, *Math. Proc. Cambridge Philos. Soc.* 133 (2002) 191–204; MR1912395 (2003e:11085).
- [88] M. B. Villarino, Mertens' proof of Mertens' theorem, math.HO/0504289.
- [89] S. R. Finch, Two asymptotic series, unpublished note (2003).
- [90] H. Cohen, High precision computation of Hardy-Littlewood constants, unpublished note (1999).

- [91] F. Saidak, An elementary proof of a theorem of Delange, *C. R. Math. Acad. Sci. Soc. Royale Canada* 24 (2002) 144–151; MR1940553 (2003k:11149).
- [92] D. A. Goldston, On the function  $S(T)$  in the theory of the Riemann zeta-function, *J. Number Theory* 27 (1987) 149–177; MR0909834 (89a:11086).
- [93] T. H. Chan, Lower order terms of the second moment of  $S(t)$ , math.NT/0411501.
- [94] A. Rényi, On the density of certain sequences of integers, *Acad. Serbe Sci. Publ. Inst. Math.* 8 (1955) 157–162; also in *Selected Papers*, v. 1, Akadémiai Kiadó, 1976, pp. 506–512; MR0076787 (17,944f).
- [95] M. Kac, A remark on the preceding paper by A. Rényi, *Acad. Serbe Sci. Publ. Inst. Math.* 8 (1955) 163–165; MR0076788 (17,944g).
- [96] K. E. Morrison, The polynomial analogue of a theorem of Rényi, *Proc. Amer. Math. Soc.* 133 (2005) 2897–2902; math.NT/0411089.
- [97] J. Grah, Comportement moyen du cardinal de certains ensembles de facteurs premiers, *Monatsh. Math.* 118 (1994) 91–109; MR1289851 (95e:11102).
- [98] J.-M. De Koninck, On a class of arithmetical functions, *Duke Math. J.* 39 (1972) 807–818; MR0311598 (47 #160).
- [99] E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, 3rd ed., v. 2, Chelsea, 1974, sects. 176–183, pp. 641–669.
- [100] E. Wirsing, Über die Zahlen, deren Primteiler einer gegebenen Menge angehören, *Arch. Math.* 7 (1956) 263–272; MR0083003 (18,642f).
- [101] P. Moree, Chebyshev’s bias for composite numbers with restricted prime divisors, *Math. Comp.* 73 (2004) 425–449; math.NT/0112100; MR2034131 (2005b:11154).
- [102] F. Ben saïd and J.-L. Nicolas, Even partition functions, *Sémin. Lothar. Combin.* 46 (2001/02) B46i; MR1921679 (2003g:11118).
- [103] F. Ben saïd and J.-L. Nicolas, Sur une application de la formule de Selberg-Delange, *Colloq. Math.* 98 (2003) 223–247; MR2033110 (2004m:11149).
- [104] L. Murata and K. Chinen, On a distribution property of the residual order of  $a \pmod p$ . II, *J. Number Theory* 105 (2004) 82–100; math.NT/0211083; MR2032443 (2005c:11118).

- [105] P. Moree, On the average number of elements in a finite field with order or index in a prescribed residue class, *Finite Fields Appl.* 10 (2004) 438–463; math.NT/0212220; MR2067608 (2005f:11219).
- [106] P. Moree, On the distribution of the order and index of  $g \bmod p$  over residue classes. I, *J. Number Theory* 114 (2005) 238–271; math.NT/0211259; MR2167970 (2006e:11152).
- [107] P. J. Stephens, An average result for Artin’s conjecture, *Mathematika* 16 (1969) 178–188; MR0498449 (58 #16565).
- [108] L. Carlitz, On a problem in additive arithmetic. II, *Quart. J. Math.* 3 (1932) 273–290.
- [109] D. R. Heath-Brown, The square sieve and consecutive square-free numbers, *Math. Annalen* 266 (1984) 251–259; MR0730168 (85h:11050).
- [110] A. E. Ingham, Some asymptotic formulae in the theory of numbers, *J. London Math. Soc.* 2 (1927) 202–208.
- [111] L. Mirsky, Summation formulae involving arithmetic functions, *Duke Math. J.* 16 (1949) 261–272; MR0030561 (11,15a).
- [112] L. Mirsky, On the frequency of pairs of square-free numbers with a given difference, *Bull. Amer. Math. Soc.* 55 (1949) 936–939; MR0031507 (11,161e).
- [113] D. Rearick, Correlation of semi-multiplicative functions, *Duke Math. J.* 33 (1966) 623–627; MR0200252 (34 #151).
- [114] F. Pappalardi, Square free values of the order function, *New York J. Math.* 9 (2003) 331–344; MR2028173 (2004i:11116).
- [115] D. R. Heath-Brown, The density of rational points on Cayley’s cubic surface, *Proc. Session in Analytic Number Theory and Diophantine Equations*, Bonn, 2002, ed. D. R. Heath-Brown and B. Z. Moroz, Bonner Math. Schriften, 2003, 1–33; math.NT/0210333; MR2075628 (2005d:14033).
- [116] P. Swinnerton-Dyer, Diophantine equations: progress and problems, *Arithmetic of Higher-Dimensional Algebraic Varieties*, Proc. 2002 Palo Alto workshop, ed. B. Poonen and Y. Tschinkel, Birkhäuser Boston, 2004, pp. 3–35; MR2028898 (2004k:11102).

- [117] J.-Y. Cai and E. Bach, On testing for zero polynomials by a set of points with bounded precision, *Theoret. Comput. Sci.* 296 (2003) 15–25; MR1965515 (2004m:68279).
- [118] L. Tóth, The probability that  $k$  positive integers are pairwise relatively prime, *Fibonacci Quart.* 40 (2002) 13–18; MR1885265 (2002k:11165).
- [119] G. Tenenbaum and J. Wu, *Exercices corrigés de théorie analytique et probabiliste des nombres*, Soc. Math. France, 1996, p. 25; MR1397501 (97h:11001).
- [120] T. Freiberg, The probability that 3 positive integers are pairwise coprime, unpublished note (2005).
- [121] P. Moree, Counting carefree couples, math.NT/0510003.
- [122] S. R. Finch, Unitarism and infinitarism, unpublished note (2004).
- [123] E. Cohen, An elementary method in the asymptotic theory of numbers, *Duke Math. J.* 28 (1961) 183–192; MR0140496 (25 #3916).
- [124] C. Pomerance, The expected number of random elements to generate a finite abelian group, *Period. Math. Hungar.* 43 (2001) 191–198; MR1830576 (2002h:20101).
- [125] R. L. Duncan, On the factorization of integers, *Proc. Amer. Math. Soc.* 25 (1970) 191–192; MR0252311 (40 #5532).
- [126] H. Z. Cao, On the average of exponents, *Northeast. Math. J.* 10 (1994) 291–296; MR1319087 (96b:11128).
- [127] K. Sinha, Average orders of certain arithmetical functions, *J. Ramanujan Math. Soc.* 21 (2006) 267–277; available online at <http://www.mast.queensu.ca/~skaneen/>.
- [128] J.-L. Nicolas, Petites valeurs de la fonction d’Euler, *J. Number Theory* 17 (1983) 375–388; MR0724536 (85h:11053).
- [129] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A002202, A014197, A058277.
- [130] K. Ford, The distribution of totients, *Ramanujan J.* 2 (1998) 67–151; available online at <http://www.math.uiuc.edu/~ford/papers.html>; MR1642874 (99m:11106).

- [131] P. Erdős and C. Pomerance, On the normal number of prime factors of  $\varphi(n)$ , *Rocky Mountain J. Math.* 15 (1985) 343–352; MR0823246 (87e:11112).
- [132] K. Ford, The number of solutions of  $\varphi(x) = m$ , *Annals of Math.* 150 (1999) 283–311; math.NT/9907204; MR1715326 (2001e:11099).
- [133] P. Erdős, C. Pomerance and E. Schmutz, Carmichael’s lambda function, *Acta Arith.* 58 (1991) 363–385; MR1121092 (92g:11093).
- [134] Yu. V. Nesterenko, Modular functions and transcendence questions (in Russian), *Mat. Sbornik*, v. 187 (1996) n. 9, 65–96; Engl. transl. in *Russian Acad. Sci. Sbornik Math.* 187 (1996) 1319–1348; MR1422383 (97m:11102).
- [135] W. Zudilin,  $\eta(q)$  and  $\eta(q)/\eta(q^2)$  are transcendental at  $q = 1/2$ , unpublished note (2004).
- [136] T. S. Bolis and A. Schinzel, Identities which imply that a ring is Boolean, *Bull. Greek Math. Soc.* 48 (2003) 1–5.
- [137] S. Ramanujan, Some formulae in the analytic theory of numbers, *Messenger of Math.* 45 (1916) 81–84; also in *Collected Papers*, ed. G. H. Hardy, P. V. Seshu Aiyar, and B. M. Wilson, Cambridge Univ. Press, 1927, pp. 133–135, 339–340; available online at <http://www.imsc.res.in/~rao/ramanujan/CamUnivCpapers/collectedright1.htm>.
- [138] M. Kühleitner and W. G. Nowak, The average number of solutions of the Diophantine equation  $U^2 + V^2 = W^3$  and related arithmetic functions, *Acta Math. Hungar.* 104 (2004) 225–240; math.NT/0307221; MR2075922 (2005d:11143).
- [139] J. M. Borwein and S. Choi, On Dirichlet series for sums of squares, *Ramanujan J.* 7 (2003) 95–127; available online at <http://eprints.cecm.sfu.ca/archive/00000142/>; MR2035795 (2005i:11123b).
- [140] M. I. Stronina, Integral points on circular cones (in Russian), *Izv. Vyssh. Uchebn. Zaved. Mat.* (1969) n. 8, 112–116; MR0248100 (40 #1354).
- [141] F. Fricker, Über die Verteilung der pythagoreischen Zahlentripel, *Arch. Math. (Basel)* 28 (1977) 491–494; MR0444590 (56 #2940).
- [142] K.-H. Fischer, Eine Bemerkung zur Verteilung der pythagoräischen Zahlentripel, *Monatsh. Math.* 87 (1979) 269–271; MR0538759 (80m:10034).

- [143] H. Jager, On the number of pairs of integers with least common multiple not exceeding  $x$ , *Nederl. Akad. Wetensch. Proc. Ser. A* 65 (1962) 346–350; *Indag. Math.* 24 (1962) 346–350; MR0148634 (26 #6141).
- [144] S. K. K. Choi, A.V. Kumchev and R. Osburn, On sums of three squares, math.NT/0502007.
- [145] R. E. Dressler, An elementary proof of a theorem of Erdős on the sum of divisors function, *J. Number Theory* 4 (1972) 532–536; MR0311556 (47 #118).
- [146] P. Erdős and J.-L. Nicolas, Grandes valeurs de fonctions liées aux diviseurs premiers consécutifs d'un entier, *Théorie des nombres*, Proc. 1987 Québec conf., ed. J.-M. De Koninck and C. Levesque, Gruyter, 1989, pp. 169–200; MR1024560 (90i:11098).
- [147] P. Sebah, Two prime difference series, unpublished note (2004).
- [148] D. A. Goldston, Y. Motohashi, J. Pintz and C. Y. Yildirim, Small gaps between primes exist, math.NT/0505300.
- [149] D. A. Goldston, S. W. Graham, J. Pintz and C.Y. Yilidirm, Small gaps between primes or almost primes, math.NT/0506067.
- [150] H. Maier and C. Pomerance, Unusually large gaps between consecutive primes, *Trans. Amer. Math. Soc.* 322 (1990) 201–237; MR0972703 (91b:11093).
- [151] J. Pintz, Very large gaps between consecutive primes, *J. Number Theory* 63 (1997) 286–301; MR1443763 (98c:11092).
- [152] P. Borwein and W. Dykshoorn, An interesting infinite product, *J. Math. Anal. Appl.* 179 (1993) 203–207; MR1244958 (94j:11131).
- [153] E. L. Basor and C. A. Tracy, The Fisher-Hartwig conjecture and generalizations, *Physica A* 177 (1991) 167–173; MR1137031 (92k:82003).
- [154] E. L. Basor and K. E. Morrison, The Fisher-Hartwig conjecture and Toeplitz eigenvalues, *Linear Algebra Appl.* 202 (1994) 129–142; available online at <http://www.calpoly.edu/~kmorriso/Research/research.html>; MR1288485 (95g:47037).
- [155] I. V. Krasovsky, Gap probability in the spectrum of random matrices and asymptotics of polynomials orthogonal on an arc of the unit circle, *Internat. Math. Res. Notices* 2004, n. 25, 1249–1272. MR2047176 (2005d:60086).

- [156] T. Ehrhardt, Dyson's constant in the asymptotics of the Fredholm determinant of the sine kernel, *Comm. Math. Phys.* 262 (2006) 317–341; math.FA/0401205; MR2200263 (2006k:82090).
- [157] P. Deift, A. Its, I. Krasovsky and X. Zhou, The Widom-Dyson constant for the gap probability in random matrix theory, math.FA/0601535.
- [158] W. Duke and Ö. Imamoglu, Special values of multiple gamma functions, *J. Théor. Nombres Bordeaux* (to appear).
- [159] K. Ford and A. Zaharescu, On the distribution of imaginary parts of zeros of the Riemann zeta function, *J. Reine Angew. Math.* 579 (2005) 145–158; math.NT/0405459; MR2124021 (2005i:11114).
- [160] D. W. Farmer, Mean values of the logarithmic derivative of the zeta function and the GUE hypothesis, math.NT/9412220.
- [161] H. L. Montgomery and U. M. A. Vorhauer, Greedy sums of distinct squares, *Math. Comp.* 73 (2004) 493–513; MR2034134 (2005b:11026).
- [162] G. H. Hardy, On certain oscillating series, *Quart. J. Pure Appl. Math.* 38 (1907) 269–288.
- [163] G. H. Hardy, *Divergent Series*, Oxford Univ. Press, 1949, p. 77; MR0030620 (11,25a).
- [164] N. D. Elkies, Puzzle involving  $x - x^2 + x^4 - x^8 + x^{16} - + \dots$ , unpublished note (2004), available online at <http://math.harvard.edu/~elkies/Misc/sol8.html>.
- [165] P. J. Grabner and H.-K. Hwang, Digital sums and divide-and-conquer recurrences: Fourier expansions and absolute convergence, *Constr. Approx.* 21 (2005) 149–179; available online at <http://algo.stat.sinica.edu.tw/hk/>; MR2107936 (2005i:11097).
- [166] L. Lhote, Modélisation et approximation de sources complexes, <http://users.info.unicaen.fr/~llhote/recherches.html>.
- [167] L. Lhote, Computation of a class of continued fraction constants, *Analytic Algorithmics and Combinatorics (ANALCO)*, Proc. 2004 New Orleans workshop, <http://www.siam.org/meetings/analco04/program.htm>.
- [168] G. Maze, Existence of a limiting distribution for the binary GCD algorithm, math.GM/0504426.

- [169] R. P. Brent, The binary Euclidean algorithm, *Millennial Perspectives in Computer Science*, Proc. 1999 Oxford-Microsoft Symp., ed. J. Davies, B. Roscoe, and J. Woodcock, Palgrave, 2000, pp. 41-53; available online at <http://www.maths.anu.edu.au/~brent/pub/pub183.html>.
- [170] B. Lindström, A remark on  $B_4$ -sequences, *J. Combin. Theory* 7 (1969) 276–277; MR0249389 (40 #2634).
- [171] A. P. Li, On  $B_3$ -sequences (in Chinese), *Acta Math. Sinica* 34 (1991) 67–71; MR1107591 (92f:11037).
- [172] S. W. Graham,  $B_h$  sequences, *Analytic Number Theory*, Proc. 1995 Allerton Park conf., v. 1, ed. B. C. Berndt, H. G. Diamond, and A. J. Hildebrand, Birkhäuser, 1996, pp. 431–449; MR1399352 (97h:11019).
- [173] M. N. Kolountzakis, Problems in the additive number theory of general sets. I, Sets with distinct sums, <http://fourier.math.uoc.gr/~mk/surveys.html>.
- [174] J. Cilleruelo, New upper bounds for finite  $B_h$  sequences, *Adv. Math.* 159 (2001) 1–17; MR1823838 (2002g:11023).
- [175] B. Green, The number of squares and  $B_h[g]$  sets, *Acta Arith.* 100 (2001) 365–390; MR1862059 (2003d:11033).
- [176] J. Cilleruelo, I. Z. Ruzsa and C. Trujillo, Upper and lower bounds for finite  $B_h[g]$  sequences, *J. Number Theory* 97 (2002) 26–34; MR1939134 (2003i:11033).
- [177] A. Plagne, A new upper bound for  $B_2[2]$  sets, *J. Combin. Theory Ser. A* 93 (2001) 378–384; MR1805304 (2001k:11035).
- [178] L. Habsieger and A. Plagne, Ensembles  $B_2[2]$ : l'état se resserre, *Integers* 2 (2002), paper A2; MR1896147 (2002m:11010).
- [179] K. O'Bryant, Sidon Sets and Beatty Sequences, Ph.D. thesis, Univ. of Illinois at Urbana-Champaign, 2002; <http://www.math.ucsd.edu/~kobryant>.
- [180] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2<sup>nd</sup> ed., rev. D. R. Heath-Brown, Oxford Univ. Press, 1986, pp. 312–327; MR0882550 (88c:11049).
- [181] K. Soundararajan, Omega results for the divisor and circle problems, *Internat. Math. Res. Notices* (2003) 1987–1998; math.NT/0302010; MR1991181 (2004f:11105).
- [182] B. Conrey, Asymptotics of  $k$ -fold convolutions of 1, unpublished note (2004).



- [183] K. Maslanka, Effective method of computing Li's coefficients and their unexpected properties, math.NT/0402168.
- [184] K. Maslanka, An explicit formula relating Stieltjes constants and Li's numbers, math.NT/0406312.
- [185] Y. V. Matiyasevich, An analytic representation for the sum of reciprocals of the nontrivial zeros of the Riemann zeta function (in Russian), *Trudy Mat. Inst. Steklov.* 163 (1984) 181–182; Engl. transl. in *Proc. Steklov Inst. Math.* 163 (1985) 181–182; MR0769884 (86d:11069).
- [186] Y. V. Matiyasevich, A relationship between certain sums over trivial and nontrivial zeros of the Riemann zeta function (in Russian); *Mat. Zametki* 45 (1989) 65–70, 142; Engl. transl. in *Math. Notes* 45 (1989) 131–135; MR1002519 (90d:11099).
- [187] Y. Matsuoka, A sequence associated with the zeros of the Riemann zeta function, *Tsukuba J. Math.* 10 (1986) 249–254; MR0868651 (87m:11084).
- [188] A. Voros, Zeta functions for the Riemann zeros, *Annales Inst. Fourier (Grenoble)* 53 (2003) 665–699; MR2008436 (2004m:11141).
- [189] P. Scholl, The thinnest densest two-dimensional packing?, <http://www.home.unix-ag.org/scholl/octagon.html>.
- [190] P. Erdős, Extremal problems in number theory, *Theory of Numbers*, ed. A. L. Whiteman, Proc. Symp. Pure Math. 8, Amer. Math. Soc., 1965, pp. 181–189; MR0174539 (30 #4740).
- [191] N. Alon and D. J. Kleitman, Sum-free subsets, *A Tribute to Paul Erdős*, ed. A. Baker, B. Bollobás and A. Hajnal, Cambridge Univ. Press, 1990, pp. 13–26; MR1117002 (92f:11020).
- [192] N. Alon and J. Spencer, *The Probabilistic Method*, Wiley, 1992, pp. 9–10; MR1140703 (93h:60002).
- [193] M. N. Kolountzakis, Some applications of probability to additive number theory and harmonic analysis, *Number Theory (New York, 1991–1995)*, Springer-Verlag, 1996, pp. 229–251; <http://fourier.math.uoc.gr/~mk/publ/>; MR1420213 (98i:11061).
- [194] M. N. Kolountzakis, Selection of a large sum-free subset in polynomial time, *Inform. Process. Lett.* 49 (1994) 255–256; <http://fourier.math.uoc.gr/~mk/publ/>; MR1266722 (95h:11143).

- [195] J. Bourgain, Estimates related to sumfree subsets of sets of integers, *Israel J. Math.* 97 (1997) 71–92; MR1441239 (97m:11026).
- [196] B. Green, The Cameron-Erdős conjecture, *Bull. London Math. Soc.* 36 (2004) 769–778; math.NT/0304058; MR2083752 (2005g:11027).
- [197] B. Green, An argument of Cameron and Erdős, unpublished note (2003), available online at <http://www.maths.bris.ac.uk/~mabjg/expos.html>.
- [198] N. J. Calkin, S. R. Finch and T. B. Flowers, Difference density and aperiodic sum-free sets, *Integers* 5 (2005) A3; MR2192081 (2006i:11024).
- [199] H. Cohn and C. Umans, A group-theoretic approach to fast matrix multiplication, *Proc. 44<sup>th</sup> Symp. on Foundations of Computer Science (FOCS)*, Cambridge, IEEE, 2003, pp. 438–449; math.GR/0307321.
- [200] H. Cohn, R. Kleinberg, B. Szegedy and C. Umans, Group-theoretic algorithms for matrix multiplication, *Proc. 46<sup>th</sup> Symp. on Foundations of Computer Science (FOCS)*, Pittsburgh, IEEE, 2005, pp. 379–388; math.GR/0511460.
- [201] R. Breusch, On the distribution of the roots of a polynomial with integral coefficients, *Proc. Amer. Math. Soc.* 2 (1951) 939–941; MR0045246 (13,552b).
- [202] E. Dunne and M. McConnell, Pianos and continued fractions, *Math. Mag.* 72 (1999) 104–115; MR1708449 (2000g:00025).
- [203] R. W. Hall and K. Josić, The mathematics of musical instruments, *Amer. Math. Monthly* 108 (2001) 347–357; MR1836944 (2002b:00017).
- [204] S. Wagon, The Collatz problem, *Math. Intelligencer* 7 (1985) 72–76; MR0769812 (86d:11103).
- [205] M. Chamberland, Una actualizacio del problema  $3x + 1$ , *Butl. Soc. Catalana Mat.* 18 (2003) 19–45; Engl. transl. available online at <http://www.math.grin.edu/~chamberl/3x.html>.
- [206] E. Roosendaal, Wagon’s constant and admissible sequences up to order 1000, unpublished note (2005).
- [207] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A100982.
- [208] T. Kotnik, Computational estimation of the order of  $\zeta(1/2 + it)$ , *Math. Comp.* 73 (2004) 949–956; available online at <http://lbk.fe.uni-lj.si/tadej.html>; MR2031417 (2004i:11098).

- [209] R. Balasubramanian and K. Ramachandra, On the frequency of Titchmarsh's phenomenon for  $\zeta(s)$ . III, *Proc. Indian Acad. Sci. Sect. A* 86 (1977) 341–351; MR0506063 (58 #21968).
- [210] M. N. Huxley, Exponential sums and the Riemann zeta function. V, *Proc. London Math. Soc.* 90 (2005) 1–41; MR2107036 (2005h:11180).
- [211] A. Granville and K. Soundararajan, Extreme values of  $|\zeta(1 + it)|$ , math.NT/0501232.
- [212] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2<sup>nd</sup> ed., rev. D. R. Heath-Brown, Oxford Univ. Press, 1986, pp. 336–347; MR0882550 (88c:11049).
- [213] R. R. Hall and G. Tenenbaum, Effective mean value estimates for complex multiplicative functions, *Math. Proc. Cambridge Philos. Soc.* 110 (1991) 337–351; MR1113432 (93e:11109).
- [214] R. R. Hall, Proof of a conjecture of Heath-Brown concerning quadratic residues, *Proc. Edinburgh Math. Soc.* 39 (1996) 581–588; MR1417699 (97m:11119).
- [215] J. S. Bradley and W. N. Everitt, On the inequality  $\|f''\|^2 \leq K\|f\|\|f^{(iv)}\|$ , *Quart. J. Math.* 25 (1974) 241–252; MR0349930 (50 #2423).
- [216] A. Russell, On a fourth-order singular integral inequality, *Proc. Royal Soc. Edinburgh* 80A (1978) 249–260; MR0516226 (80g:26022).
- [217] Vũ Q. Phóng, On inequalities for powers of linear operators and for quadratic forms, *Proc. Royal Soc. Edinburgh* 89A (1981) 25–50; MR0628126 (83g:47005).
- [218] M. J. Beynon, B. M. Brown and W. D. Evans, On an inequality of the Kolmogorov type for a second-order differential expression, *Proc. Royal Soc. London A* 442 (1993) 555–569. MR1239811 (94g:26027)
- [219] Z. Ditzian, Some remarks on inequalities of Landau and Kolmogorov, *Aequationes Math.* 12 (1975) 145–151; MR0380503 (52 #1403).
- [220] S. T. M. Ackermans, *An Asymptotic Method in the Theory of Series*, Ph.D. thesis, Technische Universiteit Eindhoven, 1964; available online at <http://www.tue.nl/bib/ftp/roeen.html>.
- [221] S. V. Parter, Extreme eigenvalues of Toeplitz forms and applications to elliptic difference equations, *Trans. Amer. Math. Soc.* 99 (1961) 153–192; MR0120492 (22 #11245).

- [222] A. Boettcher and H. Widom, From Toeplitz eigenvalues through Green's kernels to higher-order Wirtinger-Sobolev inequalities, *math.FA/0412269*.
- [223] A. Boettcher and H. Widom, On the eigenvalues of certain canonical higher-order ordinary differential operators, *math.FA/0501116*.
- [224] S. R. Finch, Bessel function zeroes, unpublished note (2003).
- [225] S. R. Finch, Nash's inequality, unpublished note (2003).
- [226] S. R. Finch, Expected lifetimes and inradii, unpublished note (2005).
- [227] S. Ahlgren, B. C. Berndt, A. J. Yee and A. Zaharescu, Integrals of Eisenstein series and derivatives of  $L$ -functions, *Int. Math. Res. Notices* (2002) 1723–1738; MR1916839 (2003f:11056).
- [228] S. R. Finch, Modular forms on  $SL_2(\mathbb{Z})$ , unpublished note (2005).
- [229] J. Boersma, Determination of the eigenvalues of some infinite matrices (in Dutch), *Liber Amicorum Jos Jansen*, ed. M. J. H. Anthonissen, Technische Universiteit Eindhoven, 2001, pp. 16–21.
- [230] G. I. Arhipov, A. A. Karacuba and V. N. Cubarikov, Trigonometric integrals (in Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 43 (1979) 971–1003, 1197; Engl. transl. in *Math. USSR Izv.* 15 (1980) 211–239; MR0552548 (81f:10050).
- [231] K. Rogers, Sharp van der Corput estimates and minimal divided differences, *Proc. Amer. Math. Soc.* 133 (2005) 3543–3550; *math.CA/0311013*.
- [232] A. Biró, On a problem of Turán concerning sums of powers of complex numbers, *Acta Math. Hungar.* 65 (1994) 209–216; MR1281430 (95d:11120).
- [233] A. Y. Cheer and D. A. Goldston, Turán's pure power sum problem, *Math. Comp.* 65 (1996) 1349–1358; MR1348041 (96j:11124).
- [234] A. Biró, An improved estimate in a power sum problem of Turán, *Indag. Math.* 11 (2000) 343–358; MR1813477 (2001j:11088).
- [235] A. Biró, An upper estimate in Turán's pure power sum problem, *Indag. Math.* 11 (2000) 499–508; MR1909813 (2003f:11140).
- [236] A. Biró, Notes on a problem of Turán, *Period. Math. Hungar.* 42 (2001) 69–76; MR1832695 (2002d:11113).

- [237] J. Andersson, On some power sum problems of Turán and Erdős, *Acta Math. Hungar.* 70 (1996) 305–316; MR1386209 (97b:11115).
- [238] J. Andersson, Explicit solutions to certain inf max problems from Turan power sum theory, math.NT/0607238.
- [239] J. Andersson, Turan’s problem 10 revisited, math.NT/0609271.
- [240] T. Erdélyi, Polynomials with Littlewood-type coefficient constraints, *Approximation Theory X: Abstract and Classical Analysis*, Proc. 2001 St. Louis conf., ed. C. K. Chui, L. L. Schumaker and J. Stöckler, Vanderbilt Univ. Press, 2002, pp. 153–196; available online at <http://www.math.tamu.edu/~terdelyi/papers-online/list.html>; MR1924857 (2003e:41008).
- [241] S. V. Konyagin, On the Littlewood problem (in Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 45 (1981) 243–265, 463; Engl. transl. in *Math. USSR-Izv.* 45 (1981) 205–225; MR0616222 (83d:10045).
- [242] O. C. McGehee, L. Pigno and B. Smith, Hardy’s inequality and the  $L^1$  norm of exponential sums, *Annals of Math.* 113 (1981) 613–618; MR0621019 (83c:43002b).
- [243] J. D. Stegeman, On the constant in the Littlewood problem, *Math. Annalen* 261 (1982) 51–54; MR0675206 (84m:42006).
- [244] K. Yabuta, A remark on the Littlewood conjecture, *Bull. Fac. Sci. Ibaraki Univ. Ser. A* 14 (1982) 19–21; MR0665949 (84b:42001).
- [245] W. Zudilin, On the unique solution of  $\sum_{k=0}^{\infty} (2k+1)^2 (-x)^{k(k+1)/2} = 0$ ,  $0 < x < 1$ , unpublished note (2004).
- [246] D. Bertrand, Theta functions and transcendence, *Ramanujan J.* 1 (1997) 339–350; MR1608721 (99b:11080).
- [247] A. P. Magnus, “One ninth” stuff, <http://www.math.ucl.ac.be/~magnus/online/>.
- [248] A. I. Aptekarev, Sharp constants for rational approximations of analytic functions (in Russian), *Mat. Sbornik*, v. 193 (2002) n. 1, 3–72; Engl. transl. in *Sbornik Math.* 193 (2002) 1–72; MR1906170 (2003g:30070).
- [249] G. P. Chistyakov, A new asymptotic expansion and asymptotically best constants in Lyapunov’s theorem. I (in Russian), *Teor. Veroyatnost. i Primenen.* 46 (2001) 326–344; Engl. transl. in *Theory Probab. Appl.* 46 (2002) 226–242; MR1968689 (2005b:60048).

- [250] G. P. Chistyakov, A new asymptotic expansion and asymptotically best constants in Lyapunov's theorem. II (in Russian), *Teor. Veroyatnost. i Primenen.* 46 (2001) 573–579; Engl. transl. in *Theory Probab. Appl.* 46 (2002) 516–522; MR1976741 (2005b:60049).
- [251] G. P. Chistyakov, A new asymptotic expansion and asymptotically best constants in Lyapunov's theorem. III (in Russian), *Teor. Veroyatnost. i Primenen.* 47 (2002) 475–497; Engl. transl. in *Theory Probab. Appl.* 47 (2003) 395–414; MR1975424 (2005b:60050).
- [252] S. R. Finch, Discrepancy and uniformity, unpublished note (2004).
- [253] S. R. Finch, Planar harmonic mappings, unpublished note (2005).
- [254] D. V. Nicolau, Definite integral and iterated function expressions for the Laplace limit, submitted (2005).
- [255] G. Tenenbaum, Crible d'Ératosthène et modèle de Kubilius, *Number Theory In Progress*, v. 2, Proc. 1997 Zakopane-Kościełisko conf., ed. K. Györy, H. Iwaniec, and J. Urbanowicz, de Gruyter, 1999, pp. 1099–1129; MR1689563 (2000g:11077).
- [256] G. V. Proskurin, The distribution of the number of vertices in the strata of a random mapping (in Russian), *Teor. Veroyatnost. i Primenen.* 18 (1973) 846–852, Engl. transl. in *Theory Probab. Appl.* 18 (1973) 803–808; MR0323608 (48 #1964).
- [257] V. F. Kolchin, *Random Mappings*, Optimization Software Inc., 1986, pp. 164–171, 177–197; MR0865130 (88a:60022).
- [258] D. J. Aldous and J. Pitman, Brownian bridge asymptotics for random mappings, *Random Structures Algorithms* 5 (1994) 487–512; MR1293075 (95k:60055).
- [259] D. J. Aldous and J. Pitman, The asymptotic distribution of the diameter of a random mapping, *C. R. Math. Acad. Sci. Paris* 334 (2002) 1021–1024; MR1913728 (2003e:60014).
- [260] V. E. Stepanov, Limit distributions of certain characteristics of random mappings (in Russian) *Teor. Veroyatnost. i Primenen.* 14 (1969) 639–653; Engl. transl. in *Theory Probab. Appl.* 14 (1969) 612–626; MR0278350 (43 #4080).
- [261] J. Blum, Enumeration of the square permutations in  $S_n$ , *J. Combin. Theory Ser. A* 17 (1974) 156–161; MR0345833 (49 #10563).

- [262] E. A. Bender, Asymptotic methods in enumeration, *SIAM Rev.* 16 (1974) 485-515; errata 18 (1976) 292; MR0376369 (51 #12545) and MR0437344 (55 #10276).
- [263] H. S. Wilf, *generatingfunctionology*, 2<sup>nd</sup> ed., Academic Press, 1994, pp. 146–150; MR1277813 (95a:05002).
- [264] N. Pouyanne, On the number of permutations admitting an  $m$ -th root, *Elec. J. Combin.* 9 (2002) R3; MR1887084 (2003a:05016).
- [265] P. Flajolet, E. Fusy, X. Gourdon, D. Panario and N. Pouyanne, A hybrid of Darboux’s method and singularity analysis in combinatorial asymptotics, *Elec. J. Combin.* 13 (2006) R103.
- [266] P. Bundschuh, Zur Note von Lehmer über eine Konstante von Erdős-Turán, *Analysis* 4 (1984) 263–266; MR0780607 (86d:40003).
- [267] L. G. Lucht, Summation of a series of Erdős-Turán, *Analysis* 6 (1986) 411–412; MR0877795 (88e:11081).
- [268] M. Klazar and F. Luca, On the maximal order of numbers in the “factorisatio numerorum” problem, math.NT/0505352.
- [269] A. Knopfmacher and H. Prodinger, On Carlitz compositions, *European J. Combin.* 19 (1998) 579–589; available online at [http://www.wits.ac.za/science/number\\_theory/apublic.htm](http://www.wits.ac.za/science/number_theory/apublic.htm); MR1637748 (99j:05006).
- [270] W. M. Y. Goh and P. Hitczenko, Average number of distinct part sizes in a random Carlitz composition, *European J. Combin.* 23 (2002) 647–657; math.CO/0110178; MR1924786 (2003i:05015).
- [271] P. Hitczenko and G. Stengle, Expected number of distinct part sizes in a random integer composition, *Combin. Probab. Comput.* 9 (2000) 519–527; math.CO/0110189; MR1816100 (2002d:05008).
- [272] A. Odlyzko and B. Richmond, On the compositions of an integer, *Combinatorial Mathematics VII*, Proc. 1979 Newcastle conf., ed. R. W. Robinson, G. W. Southern and W. D. Wallis, Lect. Notes in Math. 829, Springer-Verlag, 1980, pp. 199–210; MR0611195 (83b:05004).
- [273] P. Flajolet, X. Gourdon and P. Dumas, Mellin transforms and asymptotics: harmonic sums, *Theoret. Comput. Sci.* 144 (1995) 3–58; available

- online at <http://algo.inria.fr/flajolet/Publications/publist.html>; MR1337752 (96h:68093).
- [274] S. Heubach and T. Mansour, Enumeration of 3-letter patterns in compositions, math.CO/0603285.
- [275] S. R. Finch, Series-parallel networks, unpublished note (2003).
- [276] T. Fowler, I. Gessel, G. Labelle and P. Leroux, The specification of 2-trees, *Adv. Appl. Math.* 28 (2002) 145–168; MR1888841 (2003d:05049).
- [277] G. Labelle, C. Lamathe and P. Leroux, Labelled and unlabelled enumeration of  $k$ -gonal 2-trees, *J. Combin. Theory Ser. A* 106 (2004) 193–219; *Mathematics and Computer Science. II. Algorithms, Trees, Combinatorics and Probabilities*, Proc. 2002 Versailles conf., ed. B. Chauvin, P. Flajolet, D. Gardy and A. Mokkadem, Birkhäuser Verlag, 2002, pp. 95–109; math.CO/0312424; MR2058274 (2005a:05015).
- [278] P. Nogueira, On the combinatorics of leftist trees, *Discrete Appl. Math.* 109 (2001) 253–278; MR1818241 (2002f:05090).
- [279] S. R. Finch, Bipartite,  $k$ -colorable and  $k$ -colored graphs, unpublished note (2003).
- [280] S. R. Finch, Transitive relations, topologies and partial orders, unpublished note (2003).
- [281] G. S. Joyce and I. J. Zucker, Evaluation of the Watson integral and associated logarithmic integral for the  $d$ -dimensional hypercubic lattice, *J. Phys. A* 34 (2001) 7349–7354; MR1862771 (2002i:33018).
- [282] J. A. Fill, P. Flajolet and N. Kapur, Singularity analysis, Hadamard products, and tree recurrences, *J. Comput. Appl. Math.* 174 (2005) 271–313; math.CO/0306225; MR2106441 (2005h:05012).
- [283] A. Comtet and S. N. Majumdar, Precise asymptotics for a random walker’s maximum, *J. Stat. Mech.* (2005) P06013; cond-mat/0506195.
- [284] S. N. Majumdar, A. Comtet and R. M. Ziff, Unified solution of the expected maximum of a random walk and the discrete flux to a spherical trap, cond-mat/0509613.
- [285] R. M. Ziff, Flux to a trap, *J. Stat. Phys.* 65 (1991) 1217–1233; MR1143123 (92j:82057).



- [286] E. G. Coffman, P. Flajolet, L. Flatto and M. Hofri, The maximum of a random walk and its application to rectangle packing, *Probab. Engrg. Inform. Sci.* 12 (1998) 373–386; available online at <http://pauillac.inria.fr/algo/flajolet/Publications/>; MR1631315 (99f:60127).
- [287] I. Hueter, Proof of the conjecture that the planar self-avoiding walk has distance exponent  $3/4$ , submitted (2002), <http://faculty.baruch.cuny.edu/ihueter/research.html>.
- [288] I. Hueter, On the displacement exponent of the self-avoiding walk in three and higher dimensions, submitted (2003), <http://faculty.baruch.cuny.edu/ihueter/research.html>.
- [289] I. Hueter, Self-avoiding trails and walks and the radius of gyration exponent, submitted (2003), <http://faculty.baruch.cuny.edu/ihueter/research.html>.
- [290] T. W. Burkhardt and I. Guim, Self-avoiding walks that cross a square, *J. Phys. A* 24 (1991) L1221–L1228.
- [291] M. Bousquet-Mélou, A. J. Guttmann and I. Jensen, Self-avoiding walks crossing a square, [cond-mat/0506341](http://arxiv.org/abs/cond-mat/0506341).
- [292] B. Eisenberg and G. Stengle, Minimizing the probability of a tie for first place, *J. Math. Anal. Appl.* 198 (1996) 458–472; MR1376274 (97c:60130).
- [293] J. S. Athreya and L. M. Fidkowski, Number theory, balls in boxes, and the asymptotic uniqueness of maximal discrete order statistics, *Integers* (2000), A3; MR1759421 (2002f:60104).
- [294] S. Li and C. Pomerance, Primitive roots: a survey, *Number Theoretic Methods — Future Trends*, Proc. 2<sup>nd</sup> China-Japan Seminar on Number Theory, Iizuka, 2001, ed. by S. Kanemitsu and C. Jia, Kluwer, 2002, pp. 219–231; available online at <http://www.math.dartmouth.edu/~carlp/>; MR1974142 (2004a:11106).
- [295] B. D. McKay, Experimental asymptotics for independent vertex sets in the  $m \times n$  lattice, unpublished note (1998).
- [296] J. Tromp and G. Farneback, Combinatorics of Go, *Proc. 5<sup>th</sup> Conf. on Computer and Games*, Torino, 2006, to appear; available online at <http://homepages.cwi.nl/~tromp/go/legal.html>.
- [297] J. A. Fill, On the distribution of binary search trees under the random permutation model, *Random Structures Algorithms* 8 (1996) 1–25; MR1368848 (97f:68021).

- [298] J. A. Fill and N. Kapur, Limiting distributions for additive functionals on Catalan trees, *Theoret. Comput. Sci.* 326 (2004) 69–102; math.PR/0306226; MR2094243 (2005k:68243).
- [299] P. Flajolet and R. Sedgewick, Digital search trees revisited, *SIAM Rev.* 15 (1986) 748–767; MR0850421 (87m:68014).
- [300] N. Litvak and W. R. van Zwet, On the minimal travel time needed to collect  $n$  items on a circle, *Annals of Appl. Probab.* 14 (2004) 881–902; math.PR/0405294; MR2052907 (2004m:90006).
- [301] N. Kurokawa and M. Wakayama, On  $q$ -analogues of the Euler constant and Lerch’s limit formula, *Proc. Amer. Math. Soc.* 132 (2004) 935–943; MR2045407 (2005d:33022).
- [302] H. Prodinger, Compositions and Patricia tries: No fluctuations in the variance!, *Proc. 6<sup>th</sup> Workshop on Algorithm Engineering & Experiments and 1<sup>st</sup> Workshop on Analytic Algorithmics & Combinatorics (ALENEX/ANALC)*, New Orleans, 2004, ed. L. Arge, G. F. Italiano and R. Sedgewick, SIAM, 2004, pp. 211–215; available online at [http://math.sun.ac.za/~prodinger/abstract/abs\\_203.htm](http://math.sun.ac.za/~prodinger/abstract/abs_203.htm).
- [303] C. A. Christophi and H. M. Mahmoud, The oscillatory distribution of distances in random tries, *Annals Appl. Probab.* 15 (2005) 1536–1564; math.PR/0505259.
- [304] S. Janson, Rounding of continuous random variables and oscillatory asymptotics, *Annals of Probab.*, to appear; math.PR/0509009.
- [305] D. J. Aldous, The  $\zeta(2)$  limit in the random assignment problem, *Random Structures Algorithms* 18 (2001) 381–418; MR1839499 (2002f:60015).
- [306] R. Parviainen, Random assignment with integer costs, *Combin. Probab. Comput.* 13 (2004) 103–113; MR2034305 (2005c:90059); available online at <http://www.math.uu.se/research/pub/preprints.php>.
- [307] M. Tamaki, Minimal expected ranks for the secretary problems with uncertain selection, *Game Theory, Optimal Stopping, Probability and Statistics*, ed. F. T. Bruss and L. Le Cam, Inst. Math. Stat., 2000, pp. 127–139; MR1833856 (2002d:60033).
- [308] Z. Porosinski, The full-information best choice problem with a random number of observations, *Stochastic Process. Appl.* 24 (1987) 293–307; MR0893177 (88g:60111).

- [309] Z. Porosinski, On best choice problems having similar solutions, *Statist. Probab. Lett.* 56 (2002) 321–327; MR1892993 (2003a:62010).
- [310] J. D. Petrucci, On a best choice problem with partial information, *Annals of Statist.* 8 (1980) 1171–1174; MR0585717 (81m:62154).
- [311] T. S. Ferguson, J. P. Hardwick and M. Tamaki, Maximizing the duration of owning a relatively best object, *Strategies for Sequential Search and Selection in Real Time*, Proc. 1990 Amherst conf., ed. F. T. Bruss, T. S. Ferguson and S. M. Samuels, Amer. Math. Soc., 1992, pp. 37–57; MR1160608 (93h:60066).
- [312] S. M. Samuels, Why do these quite different best-choice problems have the same solutions? *Adv. Appl. Probab.* 36 (2004) 398–416; MR2058142 (2005f:60097).
- [313] L. A. Shepp, Explicit solutions to some problems of optimal stopping, *Annals of Math. Statist.* 40 (1969) 993–1010; MR0250415 (40 #3654).
- [314] L. H. Walker, Optimal stopping variables for Brownian motion, *Annals of Probab.* 2 (1974) 317–320; MR0397867 (53 #1723).
- [315] J. D. A. Wiseman, The expected value of  $s_n/n \approx 0.792953506407\dots$ , <http://www.jdawiseman.com/papers/easymath/coin-stopping.html>.
- [316] F. T. Bruss and G. Louchard, Optimal stopping on patterns in strings generated by independent random variables, *J. Appl. Probab.* 40 (2003) 49–72; MR1953767 (2003m:60111).
- [317] D. Assaf, L. Goldstein and E. Samuel-Cahn, Two-choice optimal stopping, *Adv. Appl. Probab.* 36 (2004) 1116–1147; math.PR/0510242; MR2119857 (2006a:60067).
- [318] U. Grimm, Improved bounds on the number of ternary square-free words, *J. Integer Seq.* 4 (2001) 01.2.7; MR1892308 (2002m:05027).
- [319] X. Sun, New lower bound on the number of ternary square-free words, *J. Integer Seq.* 6 (2003) 03.3.2.
- [320] C. Richard and U. Grimm, On the entropy and letter frequencies of ternary square-free words, *Elec. J. Combin.* 11 (2004) R14; MR2035308 (2004k:68141).
- [321] R. Kenyon, An introduction to the dimer model, math.CO/0310326.
- [322] F. D. K. Roberts, A Monte Carlo solution of a two-dimensional unstructured cluster problem, *Biometrika* 54 (1967) 625–628; MR0221728 (36 #4780).

- [323] F. D. K. Roberts and S. H. Storey, A three-dimensional cluster problem, *Biometrika* 55 (1968) 258–260; MR0225352 (37 #946).
- [324] D. F. Holcomb, M. Iwasawa and F. D. K. Roberts, Clustering of randomly placed spheres, *Biometrika* 59 (1972) 207–209.
- [325] C. Domb, A note on the series expansion method for clustering problems, *Biometrika* 59 (1972) 209–211.
- [326] L. A. Santaló, *Integral Geometry and Geometric Probability*, Addison-Wesley, 1976, p. 213; MR0433364 (55 #6340).
- [327] E. N. Gilbert, Random plane networks, *J. Soc. Indust. Appl. Math.* 9 (1961) 533–543; MR0132566 (24 #A2406).
- [328] B. Bollabás and A. Stacey, Approximate upper bounds for the critical probability of oriented percolation in two dimensions based on rapidly mixing Markov chains, *J. Appl. Probab.* 34 (1997) 859–867; MR1484020 (99j:60152).
- [329] P. Balister, B. Bollobás and M. Walters, Continuum percolation with steps in the square or the disc, *Random Structures Algorithms* 26 (2005) 392–403.
- [330] P. Winkler, Dependent percolation and colliding random walks, *Random Structures Algorithms* 16 (2000) 58–84; available online at <http://www.math.dartmouth.edu/~pw/papers/pubs.html>; MR1728353 (2001c:60160).
- [331] P. Gács, Compatible sequences and a slow Winkler percolation, *Combin. Probab. Comput.* 13 (2004) 815–856; math.PR/0011008; MR2102411 (2005j:60018).
- [332] P. Gács, Clairvoyant scheduling of random walks, *Proc. 34<sup>th</sup> ACM Symp. on Theory of Computing (STOC)*, Montreal, ACM, 2002, pp. 99–108; math.PR/0109152; MR2118617 (2005i:68004).
- [333] I. Peterson, Scheduling random walks, [http://www.maa.org/mathland/mathtrek\\_1\\_22\\_01.html](http://www.maa.org/mathland/mathtrek_1_22_01.html).
- [334] J. Tromp, Simulation and a percolation problem threshold, unpublished note (2005).
- [335] I. Jensen, Counting polyominoes: A parallel implementation for cluster computing, *Computational Science – ICCS 2003. Part III*, Proc. Melbourne/St. Petersburg conf., ed. P. M. A. Sloot, D. Abrahamson, A. V. Bogdanov, J. J. Dongarra, A. Y. Zomaya and Y. E. Gorbachev, Lect. Notes in Comp. Sci. 2659, Springer-Verlag, 2003, pp. 203–212.

- [336] G. Barequet, M. Moffie, A. Ribó and G. Rote, Counting polyominoes on twisted cylinders, submitted (2004); *Proc. 2005 European Conference on Combinatorics, Graph Theory and Applications (EuroComb '05)*, Berlin, ed. S. Felsner, DMTCS conf. vol. AE (2005), pp. 369-374; available online at [http://page.mi.fu-berlin.de/~rote/about\\_me/publications.html](http://page.mi.fu-berlin.de/~rote/about_me/publications.html).
- [337] D. Hickerson, Counting horizontally convex polyominoes, *J. Integer Seq.* 2 (1999) 99.1.8; MR1722363 (2000k:05023).
- [338] M. Kiwi, M. Loeb1 and J. Matousek, Expected length of the longest common subsequence for large alphabets, *Proc. 2004 Latin American Theoretical Informatics Conf. (LATIN)*, Buenos Aires, ed. M. Farach-Colton, Lect. Notes in Comp. Sci. 2976, Springer-Verlag, 2004, pp. 302–311; math.CO/0308234.
- [339] S. N. Majumdar and S. Nechaev, Exact asymptotic results for a model of sequence alignment, q-bio.GN/0410012.
- [340] J. Blasiak, Longest common subsequences and the Bernoulli matching model: Numerical work and analyses of the  $r$ -reach simplification, math.PR/0412375.
- [341] S. R. Finch, Hammersley's path process, unpublished note (2004).
- [342] P. Sarnak, What is...an expander? *Notices Amer. Math. Soc.* 51 (2004) 762–763.
- [343] T. Novikoff, Asymptotic behavior of the random 3-regular bipartite graph, New York Univ. honors project (2002), <http://www.math.nyu.edu/Courses/V63.0393/projects/ramanujan/ramanujan.htm>.
- [344] J. Bober, On the randomness of modular inverse mappings, New York Univ. honors project (2003), <http://www.math.nyu.edu/Courses/V63.0393/projects/modp/modp.htm>.
- [345] R. P. Stanley, Longest alternating subsequences of permutations, math.CO/0511419.
- [346] H. Widom, On the limiting distribution for the longest alternating sequence in a random permutation, math.CO/0511533.
- [347] R. P. Stanley, Increasing and decreasing subsequences of permutations and their variants, math.CO/0512035.

- [348] A. C. Kaporis, L. M. Kirousis and E. G. Lalas, The probabilistic analysis of a greedy satisfiability algorithm, *Proc. 2002 European Symp. on Algorithms (ESA)*, Rome, ed. R. Möhring and R. Raman, Lect. Notes in Comp. Sci. 2461, Springer-Verlag, 2002, pp. 574–585.
- [349] M. T. Hajiaghayi and G. B. Sorkin, The satisfiability threshold of random 3-SAT is at least 3.52, math.CO/0310193.
- [350] O. Dubois, Y. Boufkhad, and J. Mandler, Typical random 3-SAT formulae and the satisfiability threshold, *Proc. 11<sup>th</sup> ACM-SIAM Symp. on Discrete Algorithms (SODA)*, San Francisco, ACM, 2000, pp. 126–127.
- [351] S. Janson, Y. C. Stamatiou, and M. Vamvakari, Bounding the unsatisfiability threshold of random 3-SAT, *Random Structures Algorithms* 17 (2000) 103–116; erratum 18 (2001) 99–100; MR1774746 (2001c:68065) and MR1799806 (2001m:68064).
- [352] A. C. Kaporis, L. M. Kirousis, Y. C. Stamatiou, M. Vamvakari, and M. Zito, Coupon collectors,  $q$ -binomial coefficients and the unsatisfiability threshold, *Seventh Italian Conf. on Theoretical Computer Science (ICTCS)*, Proc. 2001 Torino conf., ed. A. Restivo, S. Ronchi Della Rocca, and L. Roversi, Lect. Notes in Comp. Sci. 2202, Springer-Verlag, 2001, pp. 328–338.
- [353] S.-C. Chang and R. Shrock, Some exact results for spanning trees on lattices, cond-mat/0602574.
- [354] S. Friedland and U. N. Peled, Theory of computation of multidimensional entropy with an application to the monomer-dimer problem, *Adv. Appl. Math.* 34 (2005) 486–522; math.CO/0402009.
- [355] S. Friedland, E. Krop, P. H. Lundow and K. Markstrom, Computational validations of the asymptotic matching conjectures, math.CO/0603001.
- [356] S. Friedland and L. Gurvits, Generalized Friedland-Tverberg inequality: applications and extensions, math.CO/0603410.
- [357] A. D. Sokal, Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions, *Combin. Probab. Comput.* 10 (2001) 41–77; cond-mat/9904146; MR1827809 (2002c:05078).
- [358] N. L. Biggs, R. M. Damerell and D. A. Sands, Recursive families of graphs, *J. Combin. Theory Ser. B* 12 (1972) 123–131; MR0294172 (45 #3245).

- [359] F. Brenti, G. F. Royle and D. G. Wagner, Location of zeros of chromatic and related polynomials of graphs, *Canad. J. Math.* 46 (1994) 55–80; MR1260339 (94k:05077).
- [360] G. F. Royle, Planar triangulations with real chromatic roots arbitrarily close to four, math.CO/0511304.
- [361] D. Blackwell, The square-root game, *Game Theory, Optimal Stopping, Probability and Statistics*, ed. F. T. Bruss and L. Le Cam, Inst. Math. Stat., 2000, pp. 35–37.
- [362] Z. Nehari, Some inequalities in the theory of functions, *Trans. Amer. Math. Soc.* 75 (1953) 256–286; MR0056704 (15,115c).
- [363] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A058797 (due to F. T. Adams-Watters and A. Mihailovs).
- [364] M. H. Lietzke, C. W. Nestor and J. Braun, An infinite sequence of inscribed polygons, *Amer. Math. Monthly* 66 (1959) 242–243.
- [365] C. J. Grimstone, A product of cosines, *Math. Gazette* 64 (1980) 120–121.
- [366] E. Stephen, Slowly convergent infinite products, *Math. Gazette* 79 (1995) 561–565.
- [367] J. Boersma and J. K. M. Jansen, Extending Bouwkamp’s calculation of  $\rho$ , unpublished note (1996).
- [368] D. H. Bailey, J. M. Borwein, V. Kapoor and E. Weisstein, Ten problems in experimental mathematics, *Amer. Math. Monthly*, to appear; available online at <http://crd.lbl.gov/~dhbailey/dhbpapers/>.
- [369] A. R. Kitson, The prime analog of the Kepler-Bouwkamp constant, math.HO/0608186.
- [370] W. D. Smith, Several geometric Diophantine problems: nonEuclidean Pythagorean triples, simplices with rational dihedral angles, and space-filling simplices, unpublished note (2004), <http://www.math.temple.edu/wds/homepage/diophant.abs>.
- [371] M. Chowdhury, On iterates of the chaotic logistic function  $t_{n+1} = 4t_n(1 - t_n)$ , unpublished note (2001).

- [372] T. Rivoal, Quelques propriétés du développement en cotangente continue de Lehmer, unpublished manuscript (2004).
- [373] O. R. Beaver and T. Garrity, A two-dimensional Minkowski  $\phi(x)$  function, *J. Number Theory* 107 (2004) 105–134; math.NT/0210480; MR2059953 (2005g:11125).
- [374] A. Marder, Two-dimensional analogs of the Minkowski  $\phi(x)$  function, math.NT/0405446.
- [375] G. Jackson, A very curious number, *Math. Gazette* 85 (2001) 84–86.
- [376] T. G. Feeman and O. Marrero, Very curious numbers indeed!, *Math. Gazette* 88 (2004) 98–101; available online at <http://www06.homepage.villanova.edu/timothy.feeman/publications/publist.htm>.
- [377] B. Nill, Volume and lattice points of reflexive simplices, math.AG/0412480.
- [378] A. N. W. Hone, Sigma function solution of the initial value problem for Somos 5 sequences, math.NT/0501554.
- [379] A. N. W. Hone, Elliptic curves and quadratic recurrence sequences, *Bull. London Math. Soc.* 37 (2005) 161–171; MR2119015 (2005h:11111).
- [380] A. Nijenhuis and H. S. Wilf, On a conjecture of Ryser and Minc, *Nederl. Akad. Wetensch. Proc. Ser. A* 73 (1970) 151–157; *Indag. Math.* 32 (1970) 151–157; MR0269671 (42 #4566).
- [381] S. M. Zemyan, On the zeroes of the  $N^{\text{th}}$  partial sum of the exponential series, *Amer. Math. Monthly* 112 (2005) 891–909.
- [382] H. J. H. Tuentler, On the generalized Poisson distribution, *Statist. Neerlandica* 54 (2000) 374–376; math.ST/0606238.
- [383] H. P. Boas, Majorant series, *J. Korean Math. Soc.* 37 (2000) 321–337; math.CV/0001037; MR1775963 (2001j:32001).
- [384] Ó. Martín, Look-and-say biochemistry: Exponential RNA and multistranded DNA, *Amer. Math. Monthly* 113 (2006) 289–307; available online at [http://www.uam.es/personal\\_pdi/ciencias/omartin/](http://www.uam.es/personal_pdi/ciencias/omartin/).
- [385] R. W. Barnard and J. L. Lewis, On the omitted area problem, *Michigan Math. J.* 34 (1987) 13–22; MR0873015 (87m:30035).



- [386] R. W. Barnard and K. Pearce, Rounding corners of gearlike domains and the omitted area problem, *J. Comput. Appl. Math.* 14 (1986) 217–226; *Numerical Conformal Mapping*, ed. L. N. Trefethen, North-Holland, 1986, 217–226; MR0829040 (87f:30014).
- [387] L. Banjai and L. N. Trefethen, Numerical solution of the omitted area problem of univalent function theory, *Comput. Methods Funct. Theory* 1 (2001) 259–273; available online at <http://web.comlab.ox.ac.uk/oucl/publications/natr/na-01-23.html>; MR1931615 (2003k:30014).
- [388] J. L. Lewis, On the minimum area problem, *Indiana Univ. Math. J.* 34 (1985) 631–661; MR0794580 (86i:30007).
- [389] J. Waniurski, On values omitted by convex univalent mappings, *Complex Variables Theory Appl.* 8 (1987) 173–180; MR0891759 (88e:30021).
- [390] R. W. Barnard, K. Pearce and C. Campbell, A survey of applications of the Julia variation, *Ann. Univ. Mariae Curie-Sklodowska Sect. A* 54 (2000) 1–20; available online at <http://www.math.ttu.edu/~pearce/preprint.shtml>; MR1825299 (2002a:30038).
- [391] M. Bonk and A. Eremenko, Covering properties of meromorphic functions, negative curvature and spherical geometry, *Annals of Math.* 152 (2000) 551–592; math.CV/0009251; MR1804531 (2002a:30050).
- [392] S. R. Finch, Radii in geometric function theory, unpublished note (2004).
- [393] P. Bundschuh and W. Zudilin, On theorems of Gelfond and Selberg concerning integral-valued entire functions, *J. Approx. Theory* 130 (2004) 164–178; MR2100701 (2005h:41004).
- [394] A. O. Gel'fond, Sur un théorème de M. G. Polya, *Atti Reale Accad. Naz. Lincei* 10 (1929) 569–574.
- [395] A. Selberg, Über einen Satz von A. Gelfond, *Arch. Math. Naturvid.* 44 (1941) 159–170; MR0006569 (4,6f).
- [396] C. Wang, Refining the constant in a maximum principle for the Bergman space, *Proc. Amer. Math. Soc.* 132 (2004) 853–855; MR2019965 (2004i:30017).
- [397] S. Rohde, On the theorem of Hayman and Wu, *Proc. Amer. Math. Soc.* 130 (2002) 387–394; MR1862117 (2002i:30010).

- [398] P. R. Garabedian and M. Schiffer, A coefficient inequality for schlicht functions, *Annals of Math.* 61 (1955) 116–136; MR0066457 (16,579c).
- [399] C. Pommerenke, *Boundary Behaviour of Conformal Maps*, Springer-Verlag, 1992, pp. 183–185; MR1217706 (95b:30008).
- [400] N. G. Makarov and C. Pommerenke, On coefficients, boundary size and Hölder domains, *Annales Acad. Sci. Fenn. Math.* 22 (1997) 305–312. MR1469793 (98i:30021).
- [401] A. Z. Grunspan and C. Pommerenke, The Grunsky norm and some coefficient estimates for bounded functions, *Bull. London Math. Soc.* 29 (1997) 705–712; MR1468058 (98k:30024).
- [402] H. Hedenmalm and S. Shimorin, Weighted Bergman spaces and the integral means spectrum of conformal mappings, *Duke Math. J.* 127 (2005) 341–393; math.CV/0406345.
- [403] P. Kraetzer, Experimental bounds for the universal integral means spectrum of conformal maps, *Complex Variables Theory Appl.* 31 (1996) 305–309; MR1427159 (97m:30018).
- [404] K. Baranski, A. Volberg and A. Zdunik, Brennan’s conjecture and the Mandelbrot set, *Internat. Math. Res. Notices* (1998) 589–600; available online at <http://www.mth.msu.edu/~volberg/papers/brennan/bren.html>; MR1635865 (2000a:37030).
- [405] D. Beliaev and S. Smirnov, On Littlewood’s constants, *Bull. London Math. Soc.* (to appear); available online at <http://www.math.kth.se/~beliaev/>.
- [406] D. Beliaev and S. Smirnov, Harmonic measure on fractal sets, available online at <http://www.math.kth.se/~beliaev/>.
- [407] A. E. Eremenko, Lower estimate in Littlewood’s conjecture on the mean spherical derivative of a polynomial and iteration theory, *Proc. Amer. Math. Soc.* 112 (1991) 713–715; MR1065943 (92k:30008).
- [408] I. Binder, N. Makarov and S. Smirnov, Harmonic measure and polynomial Julia sets, *Duke Math. J.* 117 (2003) 343–365; available online at <http://www.math.uiuc.edu/~ilia/papers/>; MR1971297 (2004b:37088).
- [409] I. Binder and P. W. Jones, in preparation.

- [410] C. E. Love, *Differential and Integral Calculus*, 4<sup>th</sup> ed., Macmillan, 1950, pp. 286–288.
- [411] S. Reese, A universal parabolic constant, unpublished note (2004).
- [412] S. R. Finch, Convex lattice polygons, unpublished note (2003).
- [413] J. A. Johnson, G. D. Poole and J. E. Wetzel, A small cover for convex unit arcs, *Discrete Comput. Geom.* 32 (2004) 141–147; MR2060822 (2005b:52003).
- [414] P. Doyle, Z.-X. He and B. Rodin, The asymptotic value of the circle-packing rigidity constants  $s_n$ , *Discrete Comput. Geom.* 12 (1994) 105–116; MR1280580 (95j:52034).
- [415] V. A. Zalgaller, How to get out of the woods? On a problem of Bellman (in Russian), *Matematicheskoe Prosveshchenie* 6 (1961) 191–195.
- [416] G. T. Sallee, The maximal set of constant width in a lattice, *Pacific J. Math.* 28 (1969) 669–674; MR0240724 (39 #2069).
- [417] L. Zdeborová and M. Mézard, The number of matchings in random graphs, cond-mat/0603350.
- [418] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, Springer-Verlag, 1988, pp. 1–21; MR0920369 (89a:11067), MR1194619 (93h:11069) and MR1662447 (2000b:11077).
- [419] H. Cohn and N. Elkies, New upper bounds on sphere packings. I, *Annals of Math.* 157 (2003) 689–714; math.MG/0110009; MR1973059 (2004b:11096).
- [420] G. Nebe and N. J. A. Sloane, Table of Densest Packings Presently Known, available online at <http://www.research.att.com/~njas/lattices/density.html>.
- [421] H. Cohn and A. Kumar, The densest lattice in twenty-four dimensions, *Elec. Res. Announc. Amer. Math. Soc.* 10 (2004) 58–67; math.MG/0408174; MR2075897 (2005e:11089).
- [422] H. Cohn and A. Kumar, Optimality and uniqueness of the Leech lattice among lattices, math.MG/0403263.
- [423] G. L. Watson, *Integral Quadratic Forms*, Cambridge Univ. Press, 1960, pp. 29–30; MR0118704 (22 #9475).
- [424] J. Milnor and D. Husemoller, *Symmetric Bilinear Forms*, Springer-Verlag, 1973, pp. 29–39; MR0506372 (58 #22129).

- [425] A.-M. Bergé and J. Martinet, Sur la constante d’Hermite: étude historique, *Séminaire de Théorie des Nombres, Talence, 1985–1986*, exp. 8, Univ. Bordeaux I, Talence, 1986; MR0883091 (89g:11032).
- [426] S. R. Finch, Minkowski-Siegel mass constants, unpublished note (2005).
- [427] R. Bacher, A new inequality for the Hermite constants, math.NT/0603477.
- [428] T. Kennedy, Compact packings of the plane with two sizes of discs, math.MG/0407145.
- [429] T. Kennedy, A densest compact planar packing with two sizes of discs, math.MG/0412418.
- [430] T. Kennedy, Packing the Plane with Two Sizes Of Discs, available online at <http://math.arizona.edu/~tgk/>.
- [431] H. Bieri, Die extremalen konvexen 6-Ecke der Ebene, *Elem. Math.* 16 (1961) 105–106.
- [432] M. J. Mossinghoff, A \$1 problem, *Amer. Math. Monthly* 113 (2006) 385–402; MR2225472 (2006m:51021).
- [433] L. Lovász, K. Vesztergombi, U. Wagner and E. Welzl, Convex quadrilaterals and  $k$ -sets, *Towards a Theory of Geometric Graphs*, ed. J. Pach, Amer. Math. Soc., 2004, pp. 139–148; MR2065260 (2005c:05059).
- [434] B. M. Ábrego and S. Fernández-Merchant, A lower bound for the rectilinear crossing number, *Graphs Combin.* 21 (2005) 293–300.
- [435] O. Aichholzer, F. Aurenhammer and H. Krasser, On the crossing number of complete graphs, *Computing* 76 (2006) 65–176; available online at <http://www.ist.tugraz.at/publications/oaich/>; MR2174677 (2006f:05047).
- [436] J. Balogh and G. Salazar,  $k$ -sets, convex quadrilaterals, and the rectilinear crossing number of  $K_n$ , *Discrete Comput. Geom.* 35 (2006) 671–690; available online at <http://www.math.uiuc.edu/~jobal/papers.html> .
- [437] O. Aichholzer and H. Krasser, Abstract order type extension and new results on the rectilinear crossing number, *Comput. Geom.* 36 (2007) 2–15; available online at <http://www.ist.tugraz.at/publications/oaich/> .
- [438] O. Aichholzer, J. García, D. Orden and P. Ramos, New lower bounds for the number of ( $\leq k$ )-edges and the rectilinear crossing number of  $K_n$ , math.CO/0608610.

- [439] B. Ábrego, J. Balogh, S. Fernández–Merchant, J. Leños and G. Salazar, An extended lower bound on the number of  $(\leq k)$ -edges to generalized configurations of points and the pseudolinear crossing number of  $K_n$ , available online at <http://www.ifisica.uaslp.mx/~gsalazar/> .